



Study Title: A single-blind, randomised, phase II multi-centre study to determine reactogenicity and immunogenicity of heterologous prime/boost COVID-19 vaccine schedules in adolescents (COMCOV-3)

Short Title: Comparing COVID-19 Vaccine Schedule Combinations in adolescents (Com-COV3)

Ethics Ref: 21/SC/0310

IRAS Project ID: 304450

ISRCTN Number: 12348322

EudraCT Number: 2021-004267-27

OVG Study Number: OVG 2021/04

Date and Version No: . V3.0 18-Oct-2021

Chief Investigator:	Associate Professor Matthew Snape Centre for Clinical Vaccinology and Tropical Medicine University of Oxford, Churchill Hospital, Old Road, Headington, Oxford, OX3 7LE
Investigators:	Dr Katrina Cathie Professor Paul Heath, Dr Jolanta Bernatoniene, Professor Teresa Lambe, Dr Angela Minassian, Dr Bassam Hallis, Dr Philip Connor

	Dr David Turner
Sponsor:	University of Oxford
Funder:	UK Vaccine Task Force (VTF) National Institute Health Research (NIHR) Novavax (Vaccine supply)
Chief Investigator Signature:	Matthew Snape
Statistician Signature:	Xinxue Liu, Oxford Nick Andrews, PHE

Trial Title: A single-blind, randomised, phase II multi-centre study to determine reactogenicity and immunogenicity of heterologous prime/boost COVID-19 vaccine schedules in adolescents (COMCOV-3)

EudraCT Number: 2021-004267-27

Protocol Date and Version No: V3.0 18-Oct-2021

Protocol signature page

The undersigned has read and understood the trial protocol detailed above and agrees to conduct the trial in compliance with the protocol.

Principal Investigator	Signature	Site name or ID number	Date
------------------------	-----------	---------------------------	------

Table of Contents

0. KEY TRIAL CONTACTS	8
1. CONFLICT OF INTEREST DECLARATION	10
2. LAY SUMMARY	10
3. SYNOPSIS.....	11
4. ABBREVIATIONS	14
5. BACKGROUND AND RATIONALE	17
5.1. Potential benefits.....	20
5.2. Potential risks.....	21
5.2.1. Unapproved schedules and vaccines	21
5.2.2. Associated with phlebotomy	22
5.2.3. Associated with saliva sampling.....	22
5.2.4. Associated with nasal fluid sampling	23
5.2.5. Allergic reactions.....	23
5.2.6. Behaviour change.....	23
5.2.7. Specific risks from vaccines.....	23
5.2.8. Receiving a sub-optimal vaccine schedule.....	23
5.2.9. Unwanted media attention.....	23
6. OBJECTIVES AND OUTCOME MEASURES	23
7. TRIAL DESIGN	25
7.1. Setting	25
7.2. Trial duration.....	25
7.3. Study groups	25
8. PARTICIPANT IDENTIFICATION.....	26
8.1. Trial participants	26
8.2. Inclusion criteria.....	26
8.3. Exclusion criteria	26
8.3.1. Temporary exclusion criteria	28
9. TRIAL PROCEDURES.....	28
9.1. Recruitment	28
9.1.1. Identification of volunteers.....	28
9.2. Screening and eligibility assessment.....	29
9.2.1. Initial screening	29
9.2.2. Telephone screening visit(s)	29
9.2.3. Screening during Visit 1	30
9.3. Informed consent.....	30

9.4.	Randomisation	32
9.5.	Blinding and code-breaking	32
9.6.	Visits	33
9.6.1.	Visits for participants receiving their first dose of COVID-19 vaccine in the study	33
9.6.1.1	D0: Final eligibility check, enrolment and vaccination visit	33
9.6.1.1.6	Safety pause	35
9.6.1.2	Subsequent visits	36
9.6.3	Participants under quarantine	37
9.6.4	Admission of participants to hospital with COVID-19 infection	37
9.7.	Sample handling	37
9.7.1.	Sample handling for trial purposes	37
9.7.2.	Sample handling for pregnancy testing	38
9.8.	Early discontinuation/Withdrawal of participants	38
9.9.	Definition of end of trial	39
10.	TRIAL INTERVENTIONS	39
10.1.	Investigational Medical Product(s) (IMP) Description	40
10.1.1.	VACCINE A – Pfizer BioNTech (BNT162b2)	40
10.1.2.	VACCINE B – Moderna COVID-19 vaccine	40
10.1.3.	VACCINE C – Novavax, NVXCoV2373	41
10.2.	Blinding of IMPs	41
10.3.	Storage of IMP	41
10.3.1.	Vaccine A - Pfizer BioNTech (BNT162b2)	41
10.3.2.	Vaccine B – Moderna COVID-19 vaccine	42
10.3.3.	Vaccine C – Novavax, NVXCoV2373	42
10.4.	Compliance with trial treatment	42
10.5.	Accountability of the trial treatment	42
10.6.	Concomitant medication	42
10.7.	Post-trial treatment	42
10.8.	Other treatments (non-IMPs)	43
10.9.	Other interventions	43
11.	SAFETY REPORTING	43
11.1.	Safety reporting window	43
11.2.	Adverse Event Definitions	43
11.3.	Assessment results outside of normal parameters as AEs and SAEs	44
11.3.1.	Clinical	44
11.4.	Assessment of severity	45
11.5.	Assessment of causality	46
11.6.	Procedures for reporting Adverse Events	47

11.6.1.	Solicited AEs	47
11.6.2.	Unsolicited AEs.....	47
11.7.	Reporting procedures for Serious Adverse Events	47
11.7.1.	Events exempt from immediate reporting as SAEs	48
11.8.	Expectedness	48
11.8.1.	SARs.....	48
11.8.2.	Foreseeable Adverse Reactions	49
11.9.	Adverse Events of Special Interest.....	50
11.9.1.	Disease enhancement following vaccination.....	51
11.10.	SUSAR reporting.....	52
11.11.	Development Safety Update Reports	52
11.12.	Interim reviews	52
11.13.	Safety holding rules.....	52
11.14.	Contraception and pregnancy.....	53
11.14.1.	Contraception	53
11.14.2.	Pregnancy.....	53
12.	STATISTICS.....	53
12.1.	Sample size.....	53
12.2.	Description of statistical methods	54
12.3.	Missing data	55
12.4.	Interim analyses	55
13.	DATA MANAGEMENT.....	55
13.1.	Access to Data & Data Protection	55
13.2.	Data Recording.....	56
13.3.	Record keeping	56
13.4.	Source data and Case Report Forms (CRFs).....	57
13.5.	Data Quality	58
14.	Quality Assurance Procedures	58
14.1.	Risk Assessment	58
14.2.	Monitoring	58
14.3.	Trial committees	58
14.3.1.	Trial Steering Committee	58
14.3.2.	Safety Monitoring Committee	58
14.3.3.	Study Management Committee.....	59
15.	PROTOCOL DEVIATIONS.....	59
16.	SERIOUS BREACHES.....	59
17.	ETHICAL AND REGULATORY CONSIDERATIONS	60
17.1.	Declaration of Helsinki	60

17.2.	Guidelines for Good Clinical Practice	60
17.3.	Approvals	60
17.4.	Other Ethical Considerations	60
17.5.	Reporting.....	60
17.6.	Transparency in Research	60
17.7.	Participant confidentiality.....	61
17.8.	Expenses and Benefits	61
18.	FINANCE AND INSURANCE.....	61
18.1.	Funding.....	61
18.2.	Insurance.....	61
18.3.	Contractual arrangements	61
19.	PUBLICATION POLICY	62
20.	DEVELOPMENT OF A NEW PRODUCT / PROCESS OR THE GENERATION OF INTELLECTUAL PROPERTY.....	62
21.	ARCHIVING	62
22.	REFERENCES:	62
23.	APPENDIX A: SCHEDULE OF PROCEDURES.....	65
	**Procedures performed only if prime dose administered in the study	66
24.	APPENDIX B: AMENDMENT HISTORY.....	67
25.	APPENDIX C: BLOOD SAMPLING	69
26.	APPENDIX D: DEFINITION OF MYOCARDITIS.....	70

0. KEY TRIAL CONTACTS

Chief Investigator	Matthew Snape Associate Professor in General Paediatrics and Vaccinology Centre for Clinical Vaccinology and Tropical Medicine (CCVTM) University of Oxford Churchill Hospital, Oxford OX37LE United Kingdom matthew.snape@paediatrics.ox.ac.uk
Sponsor	University of Oxford Research Governance, Ethics & Assurance, Joint Research Office Boundary Brook House Churchill Drive, Headington, Oxford, OX3 7GB, United Kingdom ctrng@admin.ox.ac.uk
Funder(s)	UK Vaccine Task Force (VTF) National Institute Health Research (NIHR) Novavax (Vaccine supply)
Clinical Trials Unit	Oxford Vaccine Group, University of Oxford, Centre for Clinical Vaccinology and Tropical Medicine (CCVTM), Churchill Hospital, Oxford, OX3 7LE, United Kingdom info@ovg.ox.ac.uk

Statisticians	<p>Xinxue Liu Oxford Vaccine Group, Centre for Clinical Vaccinology and Tropical Medicine (CCVTM), University of Oxford, Churchill Hospital, Oxford, OX3 7LE, xinxue.liu@paediatrics.ox.ac.uk</p> <p>Nick Andrews Public Health England 61 Colindale Ave, London NW9 5EQ nick.andrews@phe.gov.uk</p>										
Committees	<p>Data Safety Monitoring Board Manish Sadarangani Guido Pieleles Stephen Evans Hanna M. Nohynek Krishnan Bhaskaran</p> <p>Trial Steering Group Mary Ramsay, Public Health England Robert Read, University of Southampton Paul Turner, Imperial College London Claire Cameron, Public Health Scotland</p> <p>Study Management Group</p> <table border="0"> <tr> <td>M Snape</td><td>A Minassian</td></tr> <tr> <td>P Heath</td><td>K Cathie</td></tr> <tr> <td>J Bernatoniene</td><td>B Hallis</td></tr> <tr> <td>T Lambe</td><td>P Connor</td></tr> <tr> <td></td><td>D Taylor</td></tr> </table>	M Snape	A Minassian	P Heath	K Cathie	J Bernatoniene	B Hallis	T Lambe	P Connor		D Taylor
M Snape	A Minassian										
P Heath	K Cathie										
J Bernatoniene	B Hallis										
T Lambe	P Connor										
	D Taylor										

1. CONFLICT OF INTEREST DECLARATION

One of the COVID-19 vaccines approved for use in the current pandemic, ChAdOx1 nCoV-19, was developed as a partnership between the University of Oxford, who are sponsoring and coordinating this study, and AstraZeneca. The University of Oxford and AstraZeneca have committed to making the vaccine available on a 'not for profit' basis for the duration of the current pandemic. Both parties could potentially profit from this vaccine in the future. ChAdOx1 nCoV-19 is not included in this study, but this potential conflict is specified here for the sake of completeness.

M. Snape is an investigator on the Cov001, Cov002 and COV006 studies evaluating ChAdOx1 nCoV19. These studies are funded by NIHR and receive logistical support from AstraZeneca. M Snape is an investigator on Com-COV, funded by the NIHR. M Snape is currently, or has recently been, an investigator on studies funded +/- sponsored by vaccine manufacturers including Pfizer, GlaxoSmithKline, Janssen, MCM vaccines, Novavax and Medimmune. He receives no personal financial benefit for this work.

2. LAY SUMMARY

The successful roll-out of COVID vaccines such as 'COVID-19 mRNA Vaccine BNT162b2' and the Oxford/AstraZeneca ChAdOx1 nCoV-19 vaccine has saved approximately 60 000 lives in the UK alone up to July 2021.(1) Both of these vaccines are administered as a two-dose regimen, as is the adjuvanted protein COVID-19 vaccine from Novavax, NVXCoV2373, which is under rolling review of the MHRA at the time of writing.

The use of heterologous prime/boost (mix and match) schedules of COVID-19 vaccines has already been studied in COMCOV and COMCOV2, and a schedule with ChAdOx1 nCoV-19 as the first dose, followed by BNT162b2 as the second dose has been shown to be highly immunogenic and is now deployed routinely in non-elderly populations in Canada and many northern European countries.

Use of heterologous prime/boost schedules could also potentially be of benefit in an adolescent immunisation campaign. The potential benefits (and costs) of extending the UK COVID-19 vaccine immunisation to adolescents in the UK is under active review by the JCVI, and on 13th September 2021 it was announced that a first dose of BNT162b2 was recommended for all 12 to 17-year olds in the UK, with decisions about the timing and nature of the second dose to be decided.(2)

One concern regarding adolescent immunisation is a rare side effect of inflammation of the heart muscle (myocarditis) or membrane surrounding the heart (pericarditis) that has been observed after receipt of the second dose of BNT162b2, especially in young men.(3) Mixed schedules with alternative vaccines used for the second dose, or using half doses of COVID-19 vaccines, could be an alternative that allows protection against COVID-19 while avoiding a second full dose of BNT162b2, and could also help deploy existing stocks of vaccine as effectively as possible.

Accordingly, this study will determine the side effect profile, and the immune responses, following schedules using BNT162b2 as a first dose, and a second dose administered from 8 weeks later of either BNT162b2 (full or one third dose) Moderna COVID-19 vaccine (half dose), or NVXCoV2373 (full dose).

3. SYNOPSIS

Trial Title	A single-blind, randomised, phase II multi-centre study to determine reactogenicity and immunogenicity of heterologous prime/boost COVID-19 vaccine schedules in adolescents (COMCOV-3)
Internal ref. no. (or short title)	Comparing COVID-19 Vaccine Schedule Combinations in adolescents (Com-COV3)
Trial registration	EudraCT: 2021-004267-27 ISRCTN: 12348322
Sponsor	University of Oxford, Clinical Trials and Research Governance, Joint Research Office, Boundary Brook House Churchill Drive, Headington Oxford OX3 7GB, United Kingdom
Funder	UK Vaccine Task Force (VTF) National Institute Health Research (NIHR) Novavax (Vaccine supply)
Clinical Phase	Phase II
Trial Design	Single-blind, randomised homologous/heterologous-boost vaccine administration study

Trial Participants	Adolescents aged 12 to 16 years (inclusive)		
Sample Size	A total of 360 (+ up to an additional 10%) participants will be enrolled. A participant is considered enrolled at the time they receive a vaccine in this study		
Planned Trial Period	Approximately 12 months per participant Total trial period 1 year, 4 months		
	Objectives	Outcome Measures	Timepoint(s)
Primary	To evaluate the reactogenicity of heterologous boost at least 8 weeks following BNT162B2 prime vaccine administered to adolescents.	Solicited reactions systemic	7 days after booster immunisation
Secondary	To assess safety of heterologous boost COVID-19 vaccines	Serious adverse events Adverse events of special interest	Throughout the study
	To characterise immunogenicity of heterologous & homologous boost schedules	Anti-spike immunoglobulins	D0*, 56, 70, 84, 182, 364
		Anti-nucleocapsid immunoglobulins	D0*, 56** 182, 364
		Cellular immune responses by ELISpot	D0*, 56, 70, 182, 364
	To assess reactogenicity and safety of heterologous and homologous boost schedules of COVID-19 vaccines	Solicited local reactions	7 days after prime* and boost immunisation
		Unsolicited reactions	28 days after prime* and boost immunisation
	To evaluate immunogenicity, safety and reactogenicity of COVID-19 vaccines in participants sero-positive for SARS-CoV-2 nucleocapsid IgG at enrolment, compared with seronegative	Immunogenicity, safety & reactogenicity endpoints as outlined above	Timepoints as outlined above
Exploratory	To assess incidence of SARS-CoV-2 infection in participants receiving heterologous & homologous prime/boost	Self-reported SARS-CoV-2 infection, from community testing	Throughout the study

	schedules		
	To further characterise the blood antibody response	Neutralizing and Functional antibody assays	D0*, 56, 70, 84, 182, 364
	To characterise and compare the mucosal immune response to immunisation with homologous and heterologous COVID-19 vaccines using nasal fluid (collected using SAM-strips) and saliva samples	IgA & IgG ELISA and exploratory immunological assays in participants at selected sites	D0*, 56, 70
	To determine normal ranges of markers of cardiac muscle damage in adolescents, and to assess change post-immunisation	High sensitivity troponin and N-terminal pro B-type natriuretic peptide (NT-proBNP)	D0*, 56, 70
Intervention(s) - IMP(s)	*Only for participants receiving their first dose of COVID-19 vaccine in the study **Only for participants receiving their first dose of COVID-19 vaccine in the community		

4. ABBREVIATIONS

ADE	Antibody Dependant Enhancement
AE	Adverse event
AESI	Adverse Event of Special Interest
Anti-N IgG	Anti-nucleocapsid protein Immunoglobulin G
Anti-S IgG	Anti-spike protein Immunoglobulin G
AR	Adverse reaction
CCVTM	Centre for Clinical Vaccinology and Tropical Medicine, Oxford
CEPI	Collaboration for Epidemic Preparedness Innovations
ChAdOx1	Chimpanzee adenovirus 1
ChAdOx1-nCoV-19	Oxford/AstraZeneca COVID-19 vaccine
CI	Chief Investigator
CRF	Case Report Form
CT	Clinical Trials
CTA	Clinical Trials Authorisation
DSMB	Data Safety Monitoring Board
DSUR	Development Safety Update Report
EDC	Electronic Data Capture
ELISPOT	Enzyme-linked Immunospot
FBC	Full blood count
GCP	Good Clinical Practice
GMT	Geometric Mean Titre
GP	General Practitioner
HIV	Human Immunodeficiency virus
HRA	Health Research Authority
IB	Investigators Brochure

ICF	Informed Consent Form
IM	Intramuscular
IMP	Investigational Medicinal Product
IV	Intravenous
JCVI	Joint Committee on Vaccines and Immunisation
MHRA	Medicines and Healthcare products Regulatory Agency
mRNA	Messenger ribo-nucleic-acid
NHS	National Health Service
NIHR	National Institute for Health Research
NISEC	National Immunisation Schedule Evaluation Consortium
Novavax, NVXCoV2373	Novavax COVID-19 vaccine
NT-proBNP	N-terminal pro B-type natriuretic peptide
PBMC	Peripheral blood mononuclear cell
PCR	Polymerase chain reaction
PIMS-TS	Paediatric multisystem inflammatory syndrome temporally associated with COVID-19
Pfizer BNT162b2	Pfizer COVID-19 vaccine
qPCR	Quantitative polymerase chain reaction
RES	Research Ethics Service
PB	Post-boost
PI	Principal Investigator
PIS	Participant/ Patient Information Sheet
REC	Research Ethics Committee
RSI	Reference Safety Information
SAE	Serious Adverse Event
SAM-strips	Synthetic absorbable matrix strips
SAR	Serious Adverse Reaction

SARS-CoV2	Severe acute respiratory syndrome – coronavirus 2
SMPC	Summary of Medicinal Product Characteristics
SOP	Standard Operating Procedure
SUSAR	Suspected Unexpected Serious Adverse Reactions
TMF	Trial Master File
TSG	Trials Safety Group
µg	Microgram
VAED	Vaccine antibody enhanced disease
Vp	Viral particle
VTF	Vaccine Task Force
WHO	World Health Organisation

5. BACKGROUND AND RATIONALE

In December 2019, a cluster of patients with pneumonia of unknown cause was linked to a seafood wholesale market in Wuhan, China and were later confirmed to be infected with a novel coronavirus, known as 2019-nCoV.(4) The virus was subsequently renamed to SARS-CoV-2 because it is similar to the coronavirus responsible for severe acute respiratory syndrome (SARS-CoV), a lineage B betacoronavirus. SARS-CoV-2 shares more than 79% of its sequence with SARS-CoV, and 50% with the coronavirus responsible for Middle East respiratory syndrome (MERS-CoV), a member of the lineage C betacoronavirus.(5) COVID-19 is the infectious disease caused by SARS-CoV-2. By January 2020 there was increasing evidence of human-to-human transmission as the number of cases rapidly began to increase in China. Despite unprecedented containment measures adopted by the Chinese government, SARS-CoV-2 rapidly spread across the world. The WHO declared the COVID-19 outbreak a public health emergency of international concern on 30th January 2020. Globally, as of 13th August 2021, there have been 205,338,159 confirmed cases of COVID-19, including 4,333,094 deaths, reported to the WHO.(6)

Coronaviruses (CoVs) are spherical, enveloped, large positive-sense single-stranded RNA genomes. One-fourth of their genome is responsible for coding structural proteins, such as the spike (S) glycoprotein, envelope (E), membrane (M) and nucleocapsid (N) proteins. E, M, and N are mainly responsible for virion assembly whilst the S protein is involved in receptor binding, mediating virus entry into host cells during CoVs infection via different receptors.(7) SARS-CoV-2 belongs to the phylogenetic lineage B of the genus Betacoronavirus and it recognises the angiotensin-converting enzyme 2 (ACE2) as the entry receptor.(8) It is the seventh CoV known to cause human infections and the third known to cause severe disease after SARS-CoV and MERS-CoV.

Many social measures have been undertaken in countries across the world in order to limit the spread of the virus.(9) These have included social distancing, lockdown and mask-wearing. Currently there is no definitive treatment for COVID-19. Dexamethasone has been shown to improve mortality in those with confirmed disease and an Oxygen requirement.(10) Remdesivir, a direct anti-viral, has also been shown to reduce duration of symptoms in those who have only mild disease.(11)

Many countries have already experienced 'second' and 'third' waves of infection. On the 2nd December 2020 the MHRA granted emergency authorisation for a vaccine against COVID-19, 'COVID-19 mRNA Vaccine BNT162b2' the European Medicines Agency then granted conditional

authorisation on 21st December 2020, which is now approved for use down to the age of 12 years. This was followed by emergency authorisation of the Oxford/AstraZeneca ChAdOx1 nCoV-19 vaccine on the 29th December 2020 by the UK MHRA. The MHRA then granted emergency authorisation for the mRNA COVID-19 Vaccine Moderna on 8th January 2021. The adjuvanted protein COVID-19 vaccine from Novavax, NVXCoV2373, is under rolling review of the MHRA at the time of writing. All of these vaccines were developed for use as homologous two-dose regimens. Further vaccines using different platforms are expected to be approved for use, the majority of which are expected to be approved as two dose, homologous prime/boost schedules.

While older age groups and others at most risk of disease have been prioritised in COVID-19 immunisation campaigns, recommendations for immunisation are now being extended to adolescents in many countries including the USA, Israel and Ireland, given the high rates of infection in this age group. In the USA, as of 9th August 2021 more than 3.9 million 16 to 17 year-olds and 6.4 million 12 to 15 year-olds, have received at least one dose of COVID-19 vaccine, and 3.1 and 4.6 million (respectively) have received both doses.(12) These have almost all been with the mRNA vaccines, predominantly BNT162b2. The potential benefits (and costs) of extending the UK COVID-19 vaccine immunisation to adolescents in the UK is under active review by the JCVI, and on 13th September 2021 it was announced that a first dose of BNT162b2 was recommended for all 12 to 17 year-olds in the UK, with decisions about the timing and nature of the second dose to be decided.(2)

One concern regarding adolescent immunisation is a rare side effect of inflammation of the heart muscle (myocarditis) that has been observed after receipt of the second dose of BNT162b2, especially in young men.(3) Reports from the USA Centre for Disease Control (CDC) describe that:

'As of July 30, 2021, VAERS has received 1,249 reports of myocarditis or pericarditis among people ages 30 and younger who received COVID-19 vaccine. Most cases have been reported after mRNA COVID-19 vaccination (Pfizer-BioNTech or Moderna), particularly in male adolescents and young adults. Through follow-up, including medical record reviews, CDC and FDA have confirmed 716 reports of myocarditis or pericarditis.(13) Based on this dataset, Public Health England report that:

'Data from the USA suggests that, in males aged 12 to 17 years, 9.8 cases of myocarditis were reported per million first doses given. This rises to 67 per million after the second dose.'

Fortunately, most cases of myocarditis are benign and self-limiting, although long term effects are uncertain.(3)

The possibility of using alternatives to a full dose of BNT162b2 as the second dose of vaccine is therefore being considered, and could also help deploy available stocks of vaccine as effectively as possible. Recent data from a Phase 2/3 study which enrolled 2268 participants between 5 and 11 years of age showed that a homologous two-dose regimen of 10 µg of BNT162b2 achieved comparable immunogenicity (was non-inferior) when compared to a two-dose regimen of 30 µg administered to participants aged 16 to 25 years of age [SARS-CoV-2-neutralizing antibody geometric mean titre (GMT) 1,197.6 (95% CI 1106.1, 1296.6) in participants aged 5-11 years compared to 1146.5 (95% CI 1045.5, 1257.2) in 16-25 year olds]. These new data suggest that a lower dose (one third) in the adolescent age group may have the potential to provide an equal level of immunogenicity but with an improved reactogenicity profile (14). A potential approach is therefore to use a lower dose of an mRNA vaccine, either BNT162B2 or Moderna COVID-19 vaccine. In addition to being possibly less reactogenic than, and comparably immunogenic to, a full dose, this would also allow greater numbers to be immunised with the available supply of vaccines.

The use of heterologous prime/boost schedules of COVID-19 vaccines has been studied in COMCOV and COMCOV2, to facilitate flexible immunisation programmes. The COMCOV study showed a schedule with ChAdOx1 nCoV-19 as the first dose, followed by BNT162b2 as the second dose is highly immunogenic and is now deployed routinely in non-elderly populations in Canada and many northern European countries. However, restrictions on the use of ChAdOx1 nCoV-19 in younger adults due to concerns regarding vaccine induced thrombotic thrombocytopenia mean that this would not be a suitable option for a mixed schedule in adolescents.

Another potential option for the second dose is NVXCoV2373. While not yet licensed, a phase 3 study of this vaccine showed it to be 89.7% effective at preventing SARS-CoV-2 infection in adults, without any significant safety concerns.(15) A phase 3 study currently underway in the USA (NCT04611802) includes an adolescent cohort (N=3000), in which over 1400 participants have received NVXCoV2373 with no safety concerns raised (Novavax, personal communication). Furthermore, data generated in COMCOV2 (unpublished) has shown that a schedule employing BNT162b2 followed by NVXCoV2373 is no more reactogenic than a homologous BNT162b2/BNT162B2 schedule in participants aged over 50 years.

Accordingly, this study will determine the side effect profile, and the immune responses, following schedules using BNT162b2 as a first dose (administered in this study or in the community), and a second dose administered at least 8 weeks later of either BNT162b2 (full or one third dose), Moderna COVID-19 vaccine (half dose), or NVXCoV2373 (full dose).

Many young people in the UK have already been infected with COVID-19. It is important to understand the response to vaccination in this group. Therefore, participants enrolled in this study will include those with previous proven or suspected COVID-19.

Young people aged 12 to 17 years are now being invited to receive a dose of Pfizer-BioNTech vaccine, and many may have already received this by the time the study commences. Individuals who have received this vaccination in the community will be eligible to enrol in the study at least 8 weeks after receiving their first dose.

Table 1: Investigational medicinal product(s), summary of relevant studies (Trials for BNT162b2 and Moderna COVID-19 vaccine not shown, as they have been given approval in many countries, and millions of doses have been administered).

Country	Trial	Phase	Trials registration	Vaccine	Route	Dose	Age cohorts (years)	Number of participants
NVXCoV2373								
Australia USA	Novavax	2-part, randomized, observer-blinded, placebo-controlled, Phase 1/2 trial.	NCT04368988	NVXCoV2373	IM	0.6ml 0.5ml	18-84	1500
United Kingdom	Novavax	Phase 3, Randomised, Observer-Blinded, Placebo-Controlled	EudraCT: 2020-004123-16 NCT04583995	NVXCoV2373	IM	0.5ml	18-84	15203
South Africa	Novavax	Phase 2A/B Randomized, Observer-blinded, Placebo-controlled Study	NCT04533399	NVXCoV2373	IM	0.5ml	18-64 18-84	4422
USA Mexico	Novavax	A Phase 3, Randomized, Observer-Blinded, Placebo-Controlled Stud	NCT04611802	NVXCoV2373	IM	0.5ml	>18	30000
							12 - 17	2100

5.1. Potential benefits

A 2-dose schedule of either BNT162b2 or Moderna COVID-19 vaccine is approved for use from the age of 12 years and older in the UK, and is highly effective against COVID-19. Current JCVI recommendations for paediatric COVID-19 immunisation are:

- Two doses of COVID-19 vaccines for 12 to 17 year-olds with specific underlying health conditions that put them at increased risk of COVID-19, or for household contacts of those who are severely immunocompromised(16)
- A single dose of COVID-19 vaccines for 16 to 17 year-olds(2)

Most 12 to 16 year-olds are not eligible to receive two doses of COVID-19 vaccines, which is a potential benefit that participation in this study would provide.

The approved two dose schedule of BNT162b2 will be received by a quarter of participants in this study. All other participants will receive a single full dose of BNT162b2, which is thought to provide 80% protection against hospitalisation.(2) A second dose of the vaccines included in this study is expected to further increase protection against infection.

The age group to be enrolled in this study (12 to 16 years of age) has a lower risk of severe COVID-19 disease than older age groups. Nevertheless, they are at risk of the Paediatric multisystem inflammatory syndrome temporally associated with COVID-19 (PIMS-TS).(17) SARS-Cov-2 infection results in exclusion from school for the child, and potentially leads to a need to self-isolate for the family. The prevention of SARS-COV-2 infections in this age group is therefore of direct benefit to participants.

It is hoped that the information gained from this study will contribute to the development of a safe, effective and versatile vaccine programme against COVID-19.

5.2. Potential risks

5.2.1. Unapproved schedules and vaccines

The majority of participants will be receiving a heterologous boost, which is at variance with the current UK approved schedules. Furthermore, those randomised to receive NVXCoV2373 will be receiving a vaccine that is yet to receive MHRA emergency approval. However, NVXCoV2373 has been administered to over 1400 adolescents in an ongoing clinical trial, without raising significant safety concerns. Also, mixed BNT162b2/ NVXCoV2373 and BNT162b2/Moderna schedules have already been studied in COMCOV2 and similarly found to be well tolerated with no safety concerns.

There is the potential for the schedules with a heterologous or fractional dose boost to be less immunogenic than the approved schedule with two full doses of BNT162b2 or Moderna COVID-19 vaccine. Accordingly, adolescents eligible for two doses of a COVID-19 vaccine according to JCVI guidelines as of 3rd September 2021 will not be eligible to participate in this study, and will be advised to receive COVID-19 vaccines through the NHS. For all other participants,

participation in this study will result in them having at least one additional vaccine compared with their peers.

Therefore, participants in this study would not be disadvantaged compared to their peers.

This use of an unapproved vaccine is also relevant to the issuing of “vaccine passes” for domestic and international travel purposes; these issues and efforts to ensure participants are not disadvantaged are discussed in section 9.5.

BNT162b2 and Moderna COVID-19 vaccine have been administered to many millions of individuals (including 12 – 16 year-olds), and NVXCoV2373 has been administered to over 50000 participants in clinical trials, including over 1400 in adolescents. They therefore have a well-established safety profile. Specific concerns have been raised regarding a risk of myocarditis following a second dose of BNT162b2 in young males aged 12 to 17 years (9.8 per million first doses, rising to 67 per million after the second dose). Participants will be made aware of this at the time of enrolment and this will be specified in participant and parent information sheets. Participants will be advised to seek medical attention should they experience any of the cardinal symptoms of myocarditis (chest pain or pressure; palpitations; breathlessness after exercise, at rest or lying down; sweatiness). The diagnostic criteria for myocarditis are documented in Appendix D.(18)

5.2.2. Associated with phlebotomy

Localised bruising and discomfort can occur at the site of venepuncture. Infrequently fainting may occur. These will not be documented as AEs if they occur. Blood volumes collected will adhere to EC directive 2001/20/EC for paediatric blood volume sampling which states that ‘per individual, the trial-related blood loss (including any losses in the manoeuvre) should not exceed 3 % of the total blood volume during a period of four weeks and should not exceed 1% at any single time’. Based on these calculations, we will take 16ml per sampling timepoint from participants aged 12-16 years on each visit for immunology bloods. Some participants will turn 17 during the trial and become eligible for blood donation; r they will be asked to refrain from this for the duration of their trial involvement.

5.2.3. Associated with saliva sampling

Participants may find the saliva collection process unsavoury as it is involves drooling and spitting. This will be an optional element of the study.

5.2.4. Associated with nasal fluid sampling

Localised discomfort can occur in the nostril. Infrequently, this can result in a small amount of epistaxis, which can be controlled with pressure to the affected area. This will be an optional element of the study.

5.2.5. Allergic reactions

Allergic reactions from mild to severe may occur in response to any constituent of a medicinal product's preparation. Anaphylaxis is known to occur in approximately 2.5 to 4.7 per million recipients of mRNA COVID-19 vaccines,(19) and more generally in around 1 in 1,000,000 doses of all vaccines, but can occur in response to any vaccine or medication.(20)

5.2.6. Behaviour change

Participants might feel they can modify their COVID-19 risk behaviours on the assumption that they are protected once vaccinated. Participants will be counselled that they should continue to follow all up to date government advice in relation to COVID-19 precautions during the trial.

5.2.7. Specific risks from vaccines

Please refer to section 11.8 for full details.

5.2.8. Receiving a sub-optimal vaccine schedule

It is possible that one or more of the combinations of vaccine schedule used in this trial is, on analysis found to generate a humoral or cellular immune response lower than that of the licensed homologous BNT162b2 schedule. If this were to be the case, advice would be sought from the DSMB and TSC, who might recommend further doses of vaccine to any disadvantaged participant, if possible (either via the study or through the NHS).

5.2.9. Unwanted media attention

Trial participants can be subjected to unwanted attention from the media. They will therefore be provided with access to a document outlining some suggested media guidance.

6. OBJECTIVES AND OUTCOME MEASURES

	Objectives	Outcome Measures	Timepoint(s)
Primary	To evaluate the reactogenicity of heterologous boost at least 8 weeks following BNT162B2 prime vaccine administered to adolescents.	Solicited reactions systemic	7 days after boost immunisation

Secondary	To assess safety of heterologous boost COVID-19 vaccines	Serious adverse events Adverse events of special interest	Throughout the study
	To characterise immunogenicity of heterologous & homologous boost schedules	Anti-spike immunoglobulins	D0*, 56, 70, 84, 182, 364
		Anti-nucleocapsid immunoglobulins	D0*, 56**182, 364
		Cellular immune responses by ELISpot	D0*, 56, 70, 182, 364
	To assess reactogenicity and safety of heterologous and homologous boost schedules of COVID-19 vaccines	Solicited local reactions	7 days after prime* and boost immunisation
		Unsolicited reactions	28 days after prime* and boost immunisation
	To evaluate immunogenicity, safety & reactogenicity of COVID-19 vaccines in participants seropositive for SARS-CoV-2 nucleocapsid IgG at enrolment, compared with seronegative	Immunogenicity, safety & reactogenicity endpoints as outlined above	Timepoints as outlined above
Exploratory	To assess incidence of SARS-CoV-2 infection in participants receiving heterologous & homologous prime/boost schedules	Self-reported SARS-CoV-2 infection, from community testing	Throughout the study
	To further characterise the blood antibody response	Neutralizing and Functional antibody assays	D0*, 56, 70, 84, 182, 364
	To characterise and compare the mucosal immune response to immunisation with homologous and heterologous COVID-19 vaccines using nasal fluid (collected using SAM-strips) and saliva samples	IgA & IgG ELISA and exploratory immunological assays	D0*, 56, 70
	To determine normal ranges of markers of cardiac muscle damage in adolescents, and to assess change post-immunisation	High sensitivity troponin and N-terminal pro B-type natriuretic peptide (NT-proBNP)	D0*, 56, 70

* Only for participants receiving prime vaccination in the study

** Only for participants receiving prime vaccination in the community

7. TRIAL DESIGN

A single-blind, randomised, phase II multi-centre study to determine the reactogenicity and immunogenicity of heterologous boost COVID-19 vaccine schedules.

7.1. Setting

Multicentre study conducted through academic and NHS clinical trials sites.

7.2. Trial duration

Total duration of each participant will be 10 months post boost. The total trial period will be approximately 1 year, 5 months

7.3. Study groups

A total of 360 (+ up to an additional 10%) participants will be enrolled and randomised as below:

Group	Arm	Prime (Day 0)	Boost (day 56)
(n=360)	1	BNT162b2	BNT162b2
	2	BNT162b2	BNT162b2 (one third dose)
	3	BNT162b2	Moderna (half dose)
	4	BNT162b2	NVXCoV2373

Randomisation will be performed at the time of the second dose, and stratified by study site and anti-nucleocapsid IgG status at baseline where available.

The study will be single-blind, i.e. while staff involved in study delivery will be aware of what vaccine schedule the participant is receiving, the participant themselves will remain blinded to their boost vaccine. This blind will be maintained by applying a masking tape over the vaccine syringe as detailed in the study clinical study plan. Laboratory staff will also be blinded to the vaccine schedule received.

Participants who acquire new infection with SARS-CoV-2 (as determined by SARS-CoV-2 antigen or PCR tests) will be asked to record this in their electronic/paper diary, and will continue in the study.

8. PARTICIPANT IDENTIFICATION

8.1. Trial participants

Volunteers aged 12 to 16 years inclusive at enrolment. Comorbidities of clinical definition mild/moderate/well-controlled will be permitted. Individuals of all ethnicities will be recruited, with recruitment of those identifying as Black, Asian and Minority Ethnic particularly encouraged.

8.2. Inclusion criteria

- Parent/legal guardian/Participant is willing and able to give written informed consent for participation in the trial*
- Aged 12 to 16 years inclusive at enrolment
- In good health as determined by a trial clinician. Participants may have well controlled or mild-moderate comorbidity, as long as this would not render them eligible for 2 doses of COVID-19 vaccine (see exclusion criteria below) as part of the national roll out
- Able and willing (in the Investigator's opinion) to comply with all study requirements
- Registered with a GP, and willing to allow the investigators to discuss the participant's medical history with their General Practitioner and access all medical records when relevant to study procedures.

* Parent/legal guardian to provide informed consent for participants under the age of 16. Participants aged 16 years will be assumed to be able to provide consent for themselves, however parental support will be encouraged and investigators will reserve the right to refuse enrolment if concerns arise.

8.3. Exclusion criteria

The participant may not enter the trial if ANY of the following apply:

- Previous receipt of more than one dose of a COVID-19 vaccine, or a COVID-19 vaccine other than BNT162b2
- Belonging to a cohort advised to receive 2 doses of a COVID-19 vaccine (participants at increased risk of COVID-19, or household contacts of immunocompromised, based on JCVI and 'Green Book guidelines).
- First degree relative of a study site staff member

- Any confirmed or suspected immunosuppressive or immunodeficient state; asplenia; recurrent severe infections and use of immunosuppressant medication within the past 6 months, except topical steroids or short-term oral steroids (course lasting ≤ 14 days)
- History of anaphylaxis, allergic disease or reactions likely to be exacerbated by any component of study vaccines (e.g. hypersensitivity to the active substance or any of the SmPC/IB-listed ingredients of any study vaccine). This includes latex and polyethylene glycol/macrogol (PEG)
- Pregnancy, lactation or unwillingness to practice effective contraception from enrolment to 3 months post booster vaccination, for post-menarcheal females only
- Malignancy requiring receipt of immunosuppressive chemotherapy or radiotherapy for treatment of solid organ cancer/haematological malignancy within the 6 months prior to consent.
- Bleeding disorder (e.g. factor deficiency, coagulopathy or platelet disorder), or prior history of significant bleeding or bruising following IM injections or venepuncture
- Continuous use of anticoagulants, such as coumarins and related anticoagulants (i.e. warfarin) or novel oral anticoagulants (i.e. apixaban, rivaroxaban, dabigatran and edoxaban)
- Any serious chronic illness requiring hospital specialist supervision
- Congenital cardiovascular conditions
- Severe and/or uncontrolled respiratory disease, gastrointestinal disease, liver disease, renal disease, rheumatological disease, and neurological illness (mild/moderate well controlled comorbidities are allowed)
- History of active or previous auto-immune neurological disorders (e.g. multiple sclerosis, Guillain-Barre syndrome, transverse myelitis)
- Significant renal or hepatic impairment
- Scheduled elective surgery requiring overnight admission and/or general anaesthetic during the trial
- Insufficient level of English language to undertake all study requirements in opinion of the Investigators
- Administration of immunoglobulins and/or any blood products within the three months preceding the planned administration of the vaccines
- Participants who have participated in another research trial involving an investigational product in the past 12 weeks

- Note that a prior history of confirmed or suspected COVID-19 is NOT an exclusion criterion for this study, provided the participant otherwise satisfies the health screening criteria for the study.

8.3.1. Temporary exclusion criteria

If the volunteer has any of the following, they will not be immunised that day.

- Acute respiratory illness (moderate or severe illness with or without fever)
- Fever (oral temperature greater than 37.8°C)
- Receipt of any vaccine (licensed or investigational) within 7 days before enrolment, or intent to receive within 7 days of the COVID-19 vaccines

9. TRIAL PROCEDURES

See APPENDIX A: SCHEDULE OF PROCEDURES

9.1. Recruitment

9.1.1. Identification of volunteers

Volunteers will be recruited by methods that may include use of an advertisement +/- registration form formally approved by the ethics committee(s) and distributed or posted by means such as:

- In public places, including buses and trains, with the agreement of the owner / proprietor
- In newspapers or other literature for circulation
- On radio and/or television via announcements
- On a website or social media site operated site or sponsor or with the agreement of the owner or operator (including on-line recruitment through the Oxford Vaccine Group website)
- By e-mail distribution to a group or list only with the express agreement of the network administrator or with equivalent authorisation
- By email distribution to individuals who have already given consent to be contacted for any clinical trial at the Oxford Vaccine Centre and at trial sites including COVID-19 vaccine research
- Direct mail-out using National Health Service databases: These include the National Health Applications and Infrastructure Services (NHAIS) via a NHAIS data extract or equivalent. Initial contact to potential participants will not be made by the study team.

Instead, study invitation material will be sent out on our behalf by an external company, CFH Docmail Ltd, in order to preserve the confidentiality of potential participants. CFH Docmail Ltd is accredited as having exceeded standards under the NHS Digital Data Security and Protection Toolkit (ODS ID – 8HN70)

- Oxford Vaccine Centre databases and study site databases: We may contact individuals from databases of groups within the CCVTM (including the Oxford Vaccine Centre database) and other study sites of previous trial participants who have expressed an interest in receiving information about all future studies for which they may be eligible.
- Using local GP practices or Trusts as Participant Identification Centres (PICs)

9.2. Screening and eligibility assessment

9.2.1. Initial screening

Once parents/guardians or participants express an interest in joining the trial, they will be directed to a 2-part online screening process. The first part will assess for obvious exclusion criteria. If they pass this part, they will be asked to indicate their electronic consent to cover:

- Reporting the potential participant's medical history, including any history of SARS-CoV-2 infection identified by antigen detection, PCR testing or detection of SARS-CoV2 spike or nucleocapsid antibodies in the absence of COVID-19 vaccination
- Telephone screening visits to review their medical history (if required). Requirement to be determined by review of responses to Part 2 of online questionnaire)
- Permission to contact the participant's GP for further clarification of past medical history and vaccine status, should this be clinically indicated

Once Part 1 and consent have been completed, they may progress to Part 2. Here they will be asked to give details of any medical comorbidities or medications.

Participants without a past medical history or drug history that requires further review may be invited directly to enrolment/vaccination visits.

9.2.2. Telephone screening visit(s)

Participants for whom further clarification of eligibility is required, may be invited for telephone screening visit(s), which would then be completed by member(s) of the clinical team, based on the assessment of the part 2 responses. This will be recorded in a screening CRF. This

will reduce the amount of time participants have with the clinical team during their screening procedures, should they progress to Visit 1.

We may also contact the participant's general practitioner with the permission of the volunteer. GPs will be notified at the time of enrolment (vaccination) that the subject is taking part in the study.

Volunteers will be asked to contact the study team if there are significant changes to their health status between screening and their Visit 1.

9.2.3. Screening during Visit 1

The final eligibility assessment and D0 vaccination visit will be combined into Visit 1 (V1). See Section 9.6

9.3. Informed consent

Participants aged 16 years or over will be self-consenting as per the National Institute of Health Research guidelines. However, with the participant's permission, parents/guardians will be provided with information about the study by the trial team and via information available on the trial website, and the study team will request a parent/guardian to be at the first visit. If a participant aged 16 years declines parental notification, this will necessitate contact with their named GP to corroborate their medical history before they are enrolled. This is to safeguard vulnerable young adults.

Consent will be taken by clinical staff (registered doctor or nurse) with appropriate experience and training. Where interested participants have vulnerabilities that may impair their capacity to provide informed consent, additional input or support from the individual's parents/guardians will be sought. If there is ongoing doubt about an individual's ability to provide informed consent, then they will not be enrolled in the study.

Children/adolescents aged under 16 of years will require full parental written consent as well as signed assent from the participant themselves. Individually each participant (and their parent/guardian) will have the opportunity to question an appropriately trained and delegated researcher before signing the consent form. Participants who turn 16 years during the course of the study will be required to sign a full informed consent form at the visit occurring after their 16th birthday.

Prior to consent, the participant (and their parent/guardian) will be fully informed of all aspects of the trial, the potential risks and their obligations. The following general principles will be emphasised:

A written version and verbal explanation of the Participant Information Sheet and Informed Consent will be presented to the participant of the participant detailing:

- The exact nature of the study
- What it will involve for the participant
- The implications and constraints of the protocol
- The known side effects and any risks involved in taking part
- The sample handling protocol – participants will be informed that anonymised samples taken during the study may be shared with study collaborators
- Individual results will not be shared with participants.
- The Participant Information Sheet will be made available to the participant for an appropriate amount of time (where possible this will be a minimum of 24 hours) prior to consent being obtained. A video presentation of the Participant Information Sheet may be screened to an audience, or made available for participants to access remotely. Participants will have the opportunity to individually question an appropriately trained and delegated researcher before signing consent.
- The following general principles will be emphasised:
 - o Participation in the study is entirely voluntary
 - o Refusal to participate involves no penalty or loss of medical benefits
 - o The participant may withdraw from the study at any time
 - o The participant is free to ask questions at any time to allow him or her to understand the purpose of the study and the procedures involved
 - o The participant will be informed that they will not know whether they have received an approved COVID-19 vaccine schedule. This may have implications for any travel or other activities that may require individuals to be considered ‘fully immunised’.

Participants, like the general population, will not be exempt from following the contemporaneous government COVID-19 guidance to minimise viral transmission.

Samples taken as part of the study may be sent outside of the UK and Europe to laboratories in collaboration with the University of Oxford. These will be de-identified. Volunteers will be

asked if they consent to indefinite storage of any leftover samples for use in other ethically approved research; this will be optional.

The participant will be allowed as much time as they wish to consider the information, and the opportunity to question the Investigator, their GP or other independent parties to decide whether they will participate in the study. Written informed consent will then be obtained by means of the participant/parent/guardian dated signature, and dated signature of the person who presented and obtained the informed consent. The person obtaining the consent must be suitably qualified and experienced, and be authorised to do so by the Chief/Principal Investigator and listed on the delegation log. A copy of the signed informed consent will be given to the participant/parent/legal guardian. The original signed form will be retained at the research study site, in the CRF.

9.4. Randomisation

Computer generated randomisation list will be prepared by the study statistician. Participants will be randomised 1:1:1:1 to the four study groups.

Participants will be randomised using block randomisation. Random block sizes of 4 and 8 will be used. Randomisation will be performed at the time of the second dose, and stratified by study site and anti-nucleocapsid IgG status at baseline (where available).

9.5. Blinding and code-breaking

The study will be single-blind. Staff involved in study delivery will be aware of which vaccine the participant is receiving (arm allocation); the participant will remain blinded to their boost vaccine allocation up until the day 84 visit, at which point participants will be advised of their immunisation arm. Vaccines will be prepared out of sight of the participant and the blind will be maintained by applying a masking tape over the vaccine syringe. Laboratory staff will also be blinded to the vaccine schedule received.

If the clinical condition of a participant necessitates unblinding of the participant prior to the day 84 visit, this will be undertaken according to a trial specific working instruction and group allocation sent to the attending physician. This will be done if unblinding is thought to be relevant and likely to change clinical management.

In order not to disadvantage participants in a rapidly changing landscape of rules affecting national and international travel as well as event attendance, we will make every effort to liaise

with appropriate parties to ensure participants' vaccination status is recorded in the most suitable manner.

As of the 11th June 2021, the Department of Health and Social Care has confirmed that for any future domestic certification of vaccination status, clinical trial participants will not be disadvantaged, and will be offered an "immunised" status regardless of the licencing status or vaccine schedule received. This status would be available to trial participants while blinded, and also after unblinding even if they have received an unlicensed vaccine schedule.

9.6. Visits

The study visits and procedures will be undertaken by one of the clinical trials team. The procedures to be included in each visit are documented in the schedule of attendances (Each visit is assigned a time-point and a window period, within which the visit will be conducted. If a participant cannot attend a visit, where possible, this will be re-arranged to an in-person visit within the time window. A telephone visit may be conducted instead of the in-person visit to ascertain as much relevant information as possible if the participant is unable to attend a visit in person because of quarantine or self-isolation restrictions and the participant will be out of window if the visit is postponed.

Participants may enrol in the study either before receiving a COVID-19 vaccine (study-prime) or after receiving a first dose of Pfizer-BioNTech vaccine in the community (community-prime). The details of the D0 and D56 visits will differ for these two scenarios, as described below.

9.6.1. Visits for participants receiving their first dose of COVID-19 vaccine in the study

9.6.1.1 D0: Final eligibility check, enrolment and vaccination visit

9.6.1.1.1 Informed consent

The participant will have informed consent taken as described in Section 9.3 , before proceeding to the final eligibility check component of D0 visit. A video presentation of the aims of the study and all tests to be carried out may be screened to an audience or accessed remotely before informed consent is taken. This will be pitched at a level which should generally be comprehensible to the youngest participants. Individually, each volunteer and parent/legal guardian (if applicable) will have the opportunity to question an appropriately trained and delegated researcher before signing the assent/consent form.

9.6.1.1.2 Final eligibility check D0

During the final eligibility check component of D0 visit:

If written consent is obtained, the procedures indicated in the schedule of attendances will be undertaken including:

- Confirmation of medical history, including SARS-CoV-2 infection
- Physical examination (if required, determined according to past medical history)
- Height and weight
- Observations (temperature, heart rate)
- Urine pregnancy test in females

The eligibility of the volunteer will be reviewed by a study doctor. Decisions to exclude the volunteer from enrolling in the trial or to withdraw a volunteer from the trial will be at the discretion of the Investigator.

As per Section 8.3.1 “Temporary exclusion criteria”: If a volunteer has an acute respiratory illness (moderate or severe illness with/without fever) or a fever (oral temperature > 37.8°C) at D0/D56 visit screening, the volunteer will not be enrolled that day, but may be considered for enrolment if they recover in sufficient time.

9.6.1.1.3 Samples taken at D0

- Blood tests including:
 - COVID-19 immunogenicity bloods
 - Markers of cardiac stress
- Nasal fluid and saliva sample (both optional, and at selected sites only)

9.6.1.1.4 Vaccination at D0

Participants will be considered enrolled to the trial at the point of receipt of the first vaccine administered by the study team. All vaccines will be administered intramuscularly according to specific SOPs. The participant will stay in the trial site for observation for at least 15 minutes, in case of immediate adverse events. Participants will be issued with an NHS COVID-19 vaccination record card at the first immunisation with entry of “BNT162b2” and batch number. This will be updated at the second immunisation, but boost vaccine type or batch number will not be added as this would unblind participants; instead, “COVID-19 vaccine”, “Com-COV3 Trial” will be added. We will correspond with holders of the participant’s medical information records (e.g. GPs) to enable updating of these records.

9.6.1.1.5 eDiary

Participants/parents/guardians will be given an oral thermometer, tape measure and diary (electronic, but for those who are unable to use an electronic diary, a paper version will be

made available), with instructions on use. All participants will be given the emergency 24-hour telephone number to contact the on-call study physician if needed. Participants/parents/guardians will be instructed on how to self-assess the severity of the solicited and unsolicited AEs they are entering in the diaries. There will also be space in the diary to self-document unsolicited AEs, and whether medication was taken to relieve the symptoms. It will also log any serious medical illnesses or hospital visits which may have occurred over the entire course of the study and any diagnosis of SARS-CoV-2 infection. Given the potential for physical exercise to raise troponin and NT-proBNP independently of myocardial inflammation, there will be a space in the diary to record episodes of strenuous physical activity ("defined as anything that makes you feel really out of breath") in the period between the booster vaccine and the D70 visit. Participants/parents/guardians will be asked to report on solicited AEs for 7 days from prime and boost dose (and longer if symptoms persist at day 7, until resolution or stabilisation of symptoms) and unsolicited AEs for 28 days. The Diary will collect information on the timing and severity of the following solicited AEs:

Table 2 Solicited AEs collected on post vaccination diary cards

Local solicited AEs	Pain, Tenderness, Redness, Warmth, Itch, Swelling, Induration
Systemic solicited AEs	Fever, Feverishness, Chills, Joint pains, Muscle pains, Fatigue, Headache, Malaise, Nausea, Vomiting, Diarrhoea

Post-vaccination (7 and 28 day) diary information will be reviewed by a clinician daily, and participants may be telephoned to discuss further, should there be any clinical concerns.

The diary will contain an instruction to contact the trial team by telephone should any encounter be a hospitalisation, or if they have concerns about their health.

9.6.1.1.6 Safety pause

Given that a) the homologous BNT162b2 vaccine schedule is licensed for use in 12 to 16 year-olds, b) Data from COMCOV2 demonstrates that BNT162b2 prime followed by NVXCoV2373 boost is no more reactogenic than homologous BNT162b2/BNT162b2 and c) employing fractional dose vaccines as the boost dose are expected to be less reactogenic than full dose regimens, there will be no planned safety pause or expedited review of reactogenicity data. The Moderna COVID-19 vaccine has already been approved by the MHRA for 12 to 17 year olds and no new side effects have been identified in this age group with safety data in children similar to that in adults. As in young adults, the majority of side effects were mild to moderate and related to reactogenicity such as injection site tenderness. Therefore, there will be no

planned safety pause in place for Moderna. Should significant safety concerns arise at any point (e.g. an AESI) then the DSMB will be consulted as appropriate.

9.6.1.2 Subsequent visits

Follow-up visits will take place as per the schedule of attendances described in APPENDIX A: SCHEDULE OF PROCEDURES. Participants will be assessed for local and systemic adverse events, interim history, review of diaries (paper or electronic) and blood, saliva/ nasal fluid sampling performed at these time points as detailed in the schedule of attendances. Observations and physical exam will be performed as and when clinically indicated.

Vaccination procedures will be as for D0 visit, adapted to the specific vaccine being administered as outlined in the clinical study plan. Scheduling of the second vaccination visit (D56) should be mindful of any planned routine non-COVID-19 immunisation outside the study (e.g. seasonal influenza vaccine), to avoid administration of the study COVID-19 vaccine within a period 7 days before or after the planned routine vaccine. If necessary, in this situation scheduling of the D56 visit can be delayed beyond the specified D56 study window, but must not be brought forward to less than 8 weeks following D0.

If participants experience adverse events which the investigator (physician), CI and/or DSMB chair deem to require further close observation, the participant may be admitted to an NHS hospital for observation and further medical management under the care of the Consultant on call.

9.6.2 Visits for participants receiving their first dose of COVID-19 vaccine in the community

The screening visit for these participants will include all the procedures for informed consent and eligibility assessment described above (section 9.6.1.1.1 and 9.6.1.1.2)) including date of first dose of COVID-19 vaccination in the community.

The screening visit may be combined with the D56 visit, or be scheduled to occur separately, at some time before the D56 visit.

At the D56 visit blood samples will be taken (as will nasal fluid and saliva samples, if consent has been given for this); booster vaccine will be determined by randomisation and administered; an e-diary will be supplied. These procedures will occur in the same way as described for those participants receiving their first dose of COVID-19 vaccine in the study.

Visits after D56 (i.e. D70, D84, D182, D364), will involve the same procedures for all participants, whether or not they received their first dose of COVID-19 vaccine in the study or community. These are described in APPENDIX A: SCHEDULE OF PROCEDURES.

9.6.3 Participants under quarantine

Given the evolving epidemiological situation both globally and in the UK, should a participant be unable to attend any of their scheduled or unscheduled visits, a telephone consultation will be arranged in order to obtain core study data where possible. Participants should not attend for in-person visits if they are in their period of self-isolation/quarantine.

9.6.4 Admission of participants to hospital with COVID-19 infection

With the participant's/parent's/guardian's consent, the study team will request relevant sections of medical notes on any COVID-19 episodes resulting in hospitalisation. Any data which are relevant to assessing for disease enhancement will be collected. These are likely to include, but not limited to, information on ICU admissions, clinical parameters such as oxygen saturation, respiratory rates and vital signs, need for oxygen therapy, need for ventilatory support, imaging and blood tests results, amongst others.

9.7. Sample handling

Please refer to APPENDIX C: BLOOD SAMPLING for schedule of frequency and volume of blood sampling.

9.7.1. Sample handling for trial purposes

9.7.1.1. Immunology blood tests

Immunogenicity will be assessed by a variety of immunological assays. This will include antibodies to SARS-CoV-Spike and non-Spike antigens by ELISA, ex vivo ELISpot assays for interferon gamma secreting T cell assays, and (potentially) neutralising and other functional antibody assays. Other exploratory immunological assays including cytokine analysis and other antibody assays, DNA analysis of genetic polymorphisms potentially relevant to vaccine immunogenicity and gene expression studies amongst others may be performed at the discretion of the Investigators.

Collaboration with other specialist laboratories in the UK, Europe and outside of Europe for further exploratory tests may occur. This would involve the transfer of serum, plasma, PBMC and/or other study samples to these laboratories, but these would remain anonymised. The analyses and which laboratories carry these out will be specified in the laboratory analysis plan.

Participants will be informed that there may be leftover samples of their blood (after all testing for this study is completed), and that such samples may be stored indefinitely for possible future research (exploratory immunology), including genotypic testing of genetic polymorphisms potentially relevant to vaccine immunogenicity. Participants will be able to decide if they will permit such future use of any leftover samples. With the participant's informed consent, any leftover cells and serum/plasma will be frozen indefinitely for future analysis of COVID-19 and other coronaviruses related diseases or vaccine-related responses. If a participant elects not to permit this, all of that participant's leftover samples will be discarded at the end of the trial.

Samples that are to be stored for future research will be transferred to the OVC Biobank (REC 21/SC/0161). If the participant/parent/legal guardian (as applicable) does not consent to storage in the Oxford Vaccine Centre Biobank, then all samples will be destroyed at the end of the study.

9.7.1.2. Nasal fluid & saliva samples

An exploratory analysis of mucosal immunity will be conducted at some sites using nasal fluid collected at V1, V2 and V3 using SAM-strips (synthetic absorptive matrix) and saliva samples. Analysis will be conducted initially with IgA and IgG ELISAs, with further exploratory immunology assays conducted based on results – more detail will be included in the laboratory analysis plan. The same statements regarding collaboration, storage and use of samples as for blood in Section 9.7.1.1 apply here.

9.7.1.3. Sample handling for markers of cardiac stress

These will be centrifuged and frozen at sites before shipping to a central NHS laboratory, and tested for troponin and NT-proBNP. Residual samples will then be destroyed in accordance with standard NHS processes.

9.7.2. Sample handling for pregnancy testing

Urinary pregnancy testing: For female participants urine will be tested for beta-human chorionic gonadotrophin (β -HCG) immediately prior to vaccination. This will be a point of care test and no sample will be stored.

9.8. Early discontinuation/Withdrawal of participants

In accordance with the principles of the current revision of the Declaration of Helsinki and any other applicable regulations, a participant has the right to withdraw from the study at any time

and for any reason, and is not obliged to give his or her reasons for doing so. The Investigator may withdraw the participant at any time in the interests of the participants' health and well-being. In addition, the participant may withdraw/be withdrawn for any of the following reasons:

- Administrative decision by the Investigator
- Ineligibility (either arising during the study or retrospectively, having been overlooked at screening).
- Significant protocol deviation
- Participant non-compliance with study requirements
- An AE, which requires discontinuation of the study involvement or results in inability to continue to comply with study procedures

The reason for withdrawal will be recorded in the CRF. If withdrawal is due to an AE, appropriate follow-up visits or medical care will be arranged, with the agreement of the volunteer, until the AE has resolved, stabilised or a non-trial related causality has been assigned. The DSMB or DSMB chair may recommend withdrawal of participants.

If the participant chooses to withdraw after receipt of the first dose of vaccine, they will not need to be formally unblinded, as they will all have received BNT162b2. If they withdraw after the second dose of vaccine, they will not be unblinded, unless criteria as described in 9.5 are met. since at present there is no recommendation for routine second dose in the age group being studied.

If a participant withdraws from the study, storage of samples will continue unless the participant specifically requests otherwise. Any data collected before their withdrawal will still be used in the analysis for safety and trial integrity; if the participant requests this could be de-identified following the end of the study.

In cases of participant withdrawal, long-term safety data collection, including some procedures such as safety bloods, may continue as appropriate if participants have received any vaccine doses through the trial, unless they decline any further follow-up.

9.9. Definition of end of trial

The end of the trial is the date of the last assay conducted on the last sample collected.

10. TRIAL INTERVENTIONS

See APPENDIX A: SCHEDULE OF PROCEDURES for details

10.1. Investigational Medical Product(s) (IMP) Description

Table 3: The marketing authorisation and IMP labelling status of the vaccines:

Vaccine	UK Marketing authorisation status	IMP labelling status
Pfizer/BioNTech BNT162b2	Approved for use under a temporary authorisation of the supply of an unlicensed vaccine; regulation 174 of the Human Medicines Regulations 2012.	Not IMP labelled, product will be used as supplied by manufacturer for national supply
Moderna COVID-19 vaccine	Approved for use under a temporary authorisation of the supply of an unlicensed vaccine; regulation 174 of the Human Medicines Regulations 2012	Not IMP labelled, product will be used as supplied by manufacturer for national supply
Novavax SARS-CoV-2 rS/Matrix-M1 Adjuvant NVXCoV2373	No marketing authority or emergency use approval currently	IMP labelling required

10.1.1. VACCINE A – Pfizer BioNTech (BNT162b2)

BNT162b2 is a lipid nanoparticle-formulated, nucleoside-modified mRNA vaccine that encodes trimerised SARS-CoV-2 spike glycoprotein. BNT162b2 encodes the SARS-CoV-2 full-length spike, modified by two proline mutations to lock it in the prefusion conformation and more closely mimic the intact virus with which the elicited virus-neutralizing antibodies must interact. mRNA vaccines use the pathogen's genetic code as the vaccine; this then exploits the host cells to translate the code and then make the target spike protein. The protein then acts as an intracellular antigen to stimulate the immune response. The mRNA is then degraded within days. The vaccine RNA is formulated in lipid nanoparticles (LNPs) for more efficient delivery into cells after intramuscular injection.

10.1.1.1. Dosage, scheduling and packaging

The dose of Pfizer BioNTech COVID-19 vaccine is 30µg contained in 0.3ml of the diluted vaccine. Each pack of the Pfizer BioNTech vaccine contains 195 vials with 6 full doses per vial (975 doses per pack). It is supplied with 0.9% sodium chloride diluent for injection plastic ampoules. Participants randomised to the one third dose BNT162b2 vaccine group will receive a 0.10 ml dose.

10.1.2. VACCINE B – Moderna COVID-19 vaccine

Moderna COVID-19 vaccine contains mRNA encoding the SARS-CoV-2 spike protein. The mRNA is embedded in SM-102 lipid nanoparticles. It is produced using a cell-free in vitro transcription from the corresponding DNA templates, encoding the viral spike protein of SARS-CoV-2.

10.1.2.1

Dosage, scheduling and packaging

The full dose of Moderna COVID-19 vaccine is 100µg contained in 0.5ml vaccine. The vaccine is supplied in vials containing 10 full doses. Participants randomised to the half dose Moderna COVID-19 vaccine group will receive a 0.25 ml dose containing 50 µg.

11.1.3. VACCINE C – Novavax, NVXCoV2373

Novavax, NVXCoV2373 is a nano-particle vaccine. It is constructed from the full-length wild-type (prototype Wuhan sequence) pre-fusion trimers of SARS-CoV2 spike glycoprotein. The native protein has been modified with several substitutions to limit protease cleavage and enhance thermal stability (the putative native furin cleavage site has been modified from RRAR to QQAQ and 2 proline substitutions (positions K986P and V987P) in the HR1 domain). It has also been optimised for expression in insect (*Spodoptera frugiperda*) Sf9 cells. The recombinant S-protein genes are cloned into a baculovirus vector before being transferred into Sf9 cells. These cells then produce the protein which is extracted and purified. It is co-formulated with a saponin-based adjuvant, Matrix-M1™.

10.1.3.1. Dosage, scheduling and packaging

The dose of NVXCoV2373 is 5 µg recombinant spike protein with 50 µg Matrix-M1 adjuvant (0.5ml). The vaccine is supplied in 10 full dose vials.

10.2. Blinding of IMPs

See Section 9.5 for details.

10.3. Storage of IMP

Vaccines will be stored in accordance with manufacturers' recommendations.

All movements of the study vaccines will be documented in accordance with existing standard operating procedure (SOP). Vaccine accountability, storage, shipment and handling will be in accordance with relevant SOPs and forms. To allow for participants to receive the vaccine in a short time period, multiple clinic locations may be used. In this instance vaccines will be transported in accordance with local SOP's and approvals as required.

10.3.1. Vaccine A - Pfizer BioNTech (BNT162b2)

The Pfizer BioNTech vaccine should be stored at -70°C +/- 10°C and has shelf life of 6 months. Once thawed, the vaccine may be stored for one month (for the purposes of this study, this is considered to be 30 days) at 2 to 8°C. It should be used as soon as practically possible and within 6 hours of dilution.

10.3.2. 10.3.2. Vaccine B – Moderna COVID-19 vaccine

The Moderna COVID-19 vaccine should be stored frozen at -25°C to -15°C. At this temperature, it may be stored in the unopened vial for up to 7 months. It should be stored in the original carton to protect from light. The unopened vaccine may be stored refrigerated at 2°C to 8°C, protected from light, for maximum 30 days. Once thawed the vaccine should not be re-frozen. The unopened vaccine may be stored at 8°C to 25°C up to 24 hours after removal from refrigerated conditions.

11.3.3. Vaccine C – Novavax, NVXCoV2373

SARS-CoV-2 rS and Matrix-M1 adjuvant should be stored at 2°C to 8°C and not frozen.

10.4. Compliance with trial treatment

All vaccinations will be administered by the research team and recorded in the CRF. The study medication will be at no time in the possession of the participant and compliance will not, therefore, be an issue.

10.5. Accountability of the trial treatment

Accountability of the IMPs will be conducted in accordance with the relevant SOPs.

10.6. Concomitant medication

As set out by the exclusion criteria, volunteers may not enter the study if they have received any vaccine within 7 days before enrolment (which is defined as occurring at the time of receipt of the first study vaccine), or intend to receive within 7 days after the COVID-19 vaccine, and the subsequent COVID-19 immunisation should be scheduled to given at least 7 days before or after any non-study vaccine. Volunteers may not enter the study if they have participated in another research trial involving an investigational product in the previous 12 weeks, or if they have used immunosuppressant medication within 6 months prior to enrolment or if receipt is planned at any time during the study period (except topical steroids and short course of low dose steroids < 14 days). Concomitant medications taken at enrolment will be recorded, as will new medications within the 28 days after each immunisation.

10.7. Post-trial treatment

Decisions regarding the need for a boost dose in any of the heterologous prime/boost schedules, the nature of any boost dose and mode of delivery (e.g. NHS vs study site) will be made in consultation with the DSMB and trial steering committee.

10.8. Other treatments (non-IMPs)

Participants will be advised that they may take paracetamol prophylactically after vaccine administration. This will be from the participants own supplies rather than supplied by the study team.

10.9. Other interventions

There are no additional interventions other than those specified in this protocol.

11. SAFETY REPORTING

11.1. Safety reporting window

Safety reporting for the trial will commence once the first participant is consented; and will end 10 months after the last participant has received the second dose of an IMP for SAEs and Adverse Events of Special Interest (AESIs).

For individual participants the reporting period begins when they are consented, in person at the V1, and ends 12 months after the first dose of vaccine for SAEs and AESIs.

All adverse events (AEs) that result in a participant's withdrawal from the study will be followed up until a satisfactory resolution occurs, or until a non-study related causality is assigned (if the participant consents to this).

11.2. Adverse Event Definitions

Adverse Event (AE)	Any untoward medical occurrence in a participant to whom a medicinal product has been administered, including occurrences which are not necessarily caused by or related to that product.
Adverse Reaction (AR)	An untoward and unintended response in a participant to an investigational medicinal product which is related to any dose administered to that participant. The phrase "response to an investigational medicinal product" means that a causal relationship between a trial medication and an AE is at least a reasonable possibility, i.e. the relationship cannot be ruled out. All cases judged by either the reporting medically qualified professional or the Sponsor as having a reasonable suspected causal relationship to the trial medication qualify as adverse reactions.
Adverse Events of Special Interest (AESI)	Adverse events identified as being of particular relevance to the IMP's. These will also be reported as an SAE, if meeting SAE criteria (e.g. hospitalisation)
Serious Adverse Event	A serious adverse event is any untoward medical occurrence that:

(SAE)	<ul style="list-style-type: none"> - Results in death - Is life-threatening - Requires inpatient hospitalisation or prolongation of existing hospitalisation - Results in persistent or significant disability/incapacity - Consists of a congenital anomaly or birth defect <p>Other ‘important medical events’ may also be considered a serious adverse event when, based upon appropriate medical judgement, the event may jeopardise the participant and may require medical or surgical intervention to prevent one of the outcomes listed above.</p> <p>NOTE: The term "life-threatening" in the definition of "serious" refers to an event in which the participant was at risk of death at the time of the event; it does not refer to an event which hypothetically might have caused death if it were more severe.</p>
Serious Adverse Reaction (SAR)	An adverse event that is both serious and, in the opinion of the reporting Investigator, believed with reasonable probability to be due to one of the trial treatments, based on the information provided.
Suspected Unexpected Serious Adverse Reaction (SUSAR)	<p>A serious adverse reaction, the nature and severity of which is not consistent with the Reference Safety Information for the medicinal product in question set out:</p> <ol style="list-style-type: none"> 1. In the case of a product with a marketing authorisation, in the approved summary of product characteristics (SmPC) for that product 2. In the case of any other investigational medicinal product, in the approved investigator’s brochure (IB) relating to the trial in question

NB: to avoid confusion or misunderstanding of the difference between the terms “serious” and “severe”, the following note of clarification is provided: “Severe” is often used to describe intensity of a specific event, which may be of relatively minor medical significance. “Seriousness” is the regulatory definition supplied above.

11.3. Assessment results outside of normal parameters as AEs and SAEs

11.3.1. Clinical

Abnormal clinical findings from medical history or examination will be assessed as to their clinical significance throughout the trial. If an abnormal finding is deemed to be clinically

significant, the participant will be informed and appropriate medical care arranged with the permission of the participant as per Section 9.5.

11.4. Assessment of severity

The severity of clinical and laboratory adverse events will be assessed according to scales listed in the Clinical Study Plan and in Table 4 below.

Table 4: Severity grading for local adverse events

Adverse Event	Grade	Intensity
Pain at injection site	1	Pain that is easily tolerated
	2	Pain that interferes with daily activity
	3	Pain that prevents daily activity
	4	A&E visit or hospitalization
Tenderness	1	Mild discomfort to touch
	2	Discomfort with movement
	3	Significant discomfort at rest
	4	A&E visit or hospitalization
Erythema at injection site*	1	2.5 - 5 cm
	2	5.1 - 10 cm
	3	>10 cm
	4	Necrosis or exfoliative dermatitis
Induration/Swelling at injection site	1	2.5 – 5 cm and does not interfere with activity
	2	5.1 - 10 cm or interferes with activity
	3	>10 cm or prevents daily activity
	4	Necrosis
*erythema \leq2.5cm is an expected consequence of skin puncture and will therefore not be considered an adverse event		

Table 6: Severity grading for local and systemic AEs

GRADE 0	None
GRADE 1	Mild: Transient or mild discomfort (< 48 hours); No interference with activity; No medical intervention/therapy required
GRADE 2	Moderate: Mild to moderate limitation in activity – some assistance may be needed; no or minimal medical intervention/therapy required
GRADE 3	Severe: Marked limitation in activity, some assistance usually required; medical intervention/therapy required.
GRADE 4	Potentially Life-threatening: Requires assessment in A&E or hospitalisation

11.5. Assessment of causality

For every recorded AE, an assessment of the relationship of the event to the administration of the vaccine will be undertaken by the CI-delegated clinician. An interpretation of the causal relationship of the intervention to the AE in question will be made, based on the type of event; the relationship of the event to the time of vaccine administration; and the known biology of the vaccine therapy. Alternative causes of the AE, such as the natural history of pre-existing medical conditions, concomitant therapy, other risk factors and the temporal relationship of the event to vaccination will be considered and investigated. Causality of SAEs will be assigned by the reporting investigator at the time of reporting, as described in SOP OVC005 Safety Reporting for CTIMPs. For all other AE's causality assessment will take place during planned safety reviews, interim analyses (including if the study is paused by the DSMB due to safety concerns) and at the final safety analysis. Causality assessment will be recorded on the eCRF.

Table 5: Guidelines for assessing the relationship of vaccine administration to an AE.

0	No relationship	No temporal relationship to study product and Alternate aetiology (clinical state, environmental or other interventions); and Does not follow known pattern of response to study product
1	Unlikely	Unlikely temporal relationship to study product and Alternate aetiology likely (clinical state, environmental or other interventions) and Does not follow known typical or plausible pattern of response to study product.
2	Possible	Reasonable temporal relationship to study product; or Event not readily produced by clinical state, environmental or other

		interventions; or Similar pattern of response to that seen with other vaccines
3	Probable	Reasonable temporal relationship to study product; and Event not readily produced by clinical state, environment, or other interventions or Known pattern of response seen with other vaccines
4	Definite	Reasonable temporal relationship to study product; and Event not readily produced by clinical state, environment, or other interventions; and Known pattern of response seen with other vaccines

11.6. Procedures for reporting Adverse Events

11.6.1. Solicited AEs

Participants will be asked to record local and systemic AEs for 7 days (and longer if symptoms persist at day seven, until resolution or stabilisation) following vaccination in the electronic/paper diary (solicited AEs).

11.6.2. Unsolicited AEs

All local and systemic AEs occurring in the 28 days following each vaccination observed by the Investigator or reported by the participant, whether or not attributed to study medication, will be recorded in electronic/paper diaries or study database. All AEs that result in a participant's withdrawal from the study will be followed up until a satisfactory resolution occurs, or until a non-study related causality is assigned (if the participant consents to this) as per Section 9.8.

SAEs and AESIs will be actively solicited at each study visit throughout the entire trial period.

11.7. Reporting procedures for Serious Adverse Events

In order to comply with current regulations on SAE reporting to regulatory authorities, the event will be documented accurately and notification deadlines respected. SAEs will be reported to members of the study team immediately the Investigators become aware of their occurrence, as described in the clinical study plan. Copies of all reports will be forwarded for review to the Chief Investigator (as the Sponsor's representative) within 24 hours of the Investigator being aware of the suspected SAE. The DSMB will be notified of SAEs that are deemed possibly, probably or definitely related to study interventions; the chair of DSMB will be notified immediately (within 24 hours) of the sponsor being aware of their occurrence.

SAE/AESIs will not normally be reported immediately to the ethical committee(s) unless there is

a clinically important increase in occurrence rate, an unexpected outcome, or a new event that is likely to affect safety of trial participants, at the discretion of the Chief Investigator and/or DSMB. In addition to the expedited reporting above, the Investigator shall include all SAE/AESIs in the annual Development Safety Update Report (DSUR) report provided for COM-COV, COM-COV2 and COM-COV3.

In participants who have received the NVXCoV2373 vaccine in this study, SAE's will be reported to Novavax according to the conditions and timelines outlined in the contemporaneous version of the Clinical Trial Agreement between Novavax and the University of Oxford University for conduct of this study.

11.7.1. Events exempt from immediate reporting as SAEs

Hospitalisation for a pre-existing condition, including elective procedures planned prior to study entry, which has not worsened, does not constitute a serious adverse event. A&E attendances should not routinely be reported as SAEs unless they meet the SAE definition described above.

11.8. Expectedness

11.8.1. SARs

11.8.1.1. Pfizer BioNTech (BNT162b2)

If an SAE is considered as being an SAR to BNT162b2, section 4.8 of the BNT162b2 Summary of Product Characteristics will be used as the reference safety information to determine expectedness. Potential SARs considered to be expected include:

- Anaphylaxis
- Myocarditis/pericarditis
- Lymphadenopathy
- Facial nerve palsy

11.8.1.2. Moderna COVID-19 vaccine

If an SAE is considered as being an SAR to Moderna COVID-19 vaccine, section 4.8 of the Summary of Product Characteristics will be used as the reference safety information to determine expectedness. Potential SARs considered to be expected include:

- Anaphylaxis
- Myocarditis/pericarditis
- Lymphadenopathy
- Facial nerve palsy

11.8.1.3.

Novavax, NVXCoV2373

No SARs expected

11.8.2. Foreseeable Adverse Reactions

The foreseeable ARs following vaccination are as follows:

11.8.2.1. Pfizer BioNTech (BNT162b2)

<i>Very common</i>	<i>Common</i>	<i>Uncommon</i>	<i>Rare</i>	<i>Unknown</i>
Headache	Injection site redness	Lymphadenopathy	Acute peripheral facial paralysis	Anaphylaxis
Arthralgia	Nausea	Insomnia		Hypersensitivity
Myalgia		Pain in extremity		Myocarditis
Injection site pain/swelling		Malaise		Pericarditis
Fatigue		Injection site pruritus		
Chills				
Pyrexia				

11.8.2.2. Moderna COVID-19 vaccine

<i>Very common</i>	<i>Common</i>	<i>Uncommon</i>	<i>Rare</i>	<i>Unknown</i>
Swelling/tenderness under arm	Injection site redness, rash or hives	Injection site itching	Acute peripheral facial paralysis	Anaphylaxis
Headache	Rash		Facial swelling	Hypersensitivity
Nausea or vomiting			Dizziness	Myocarditis
Muscle/joint ache			Decreased sense of touch	Pericarditis
Injection site pain/swelling				
Tiredness				
Chills				
Fever				

11.8.2.3. Novavax, NVXCoV2373

(From the IB report of data from clinical trials)

- Reactogenicity is generally mild, and vaccinations were well tolerated.

- Following first vaccination, local reactogenicity is more frequent for NVXCoV2373 than the unadjuvanted or placebo regimens.
- Tenderness and pain were the most frequent local AEs.
- Systemic reactogenicity were individually less frequent but were observed with greater frequency in the SARS-CoV-2 rS/Matrix-M1 adjuvant groups.
- Headache, fatigue, and myalgia were the most frequent systemic AEs.
- Following second vaccination, NVXCoV2373 induced greater local and systemic reactogenicity, but the majority of reported symptoms remained at grade ≤ 1 .
- Mean duration of reactogenicity events was ≤ 2 days without appreciable change in duration with second vaccination.
- Severe reactogenicity was infrequent (2 events after Dose 1 and 8 events after Dose 2), occurring more often with second vaccination and for systemic events, with placebo subjects citing similar frequencies as those receiving SARS-CoV-2rS.
- No subjects sought medical intervention or refused second vaccination because of reactogenicity.

11.9. Adverse Events of Special Interest

Table 6: AESIs

Immunologic	Anaphylaxis	
Neurological	Isolated anosmia/ageusia* Guillain-Barre Syndrome Acute disseminated encephalomyelitis (ADEM) Aseptic meningitis	Meningoencephalitis Peripheral facial nerve palsy Generalised convulsion Myelitis
Haematological	Thrombosis** Stroke Thrombocytopenia*** Eosinophilia****	Coagulation disorder (includes coagulopathy, thrombosis, thromboembolism, internal/external bleed and stroke)
Cardiac	Acute cardiovascular injury (includes myocarditis#, pericarditis#, arrhythmias, heart failure, infarction)	
Dermatological	Chilblain-like lesions Single organ cutaneous vasculitis	Erythema multiforme Alopecia
Gastrointestinal	Acute liver injury †††	Appendicitis

Respiratory	ARDS††
Renal	Acute kidney injury
Other	COVID-19 disease† SARS-CoV2 positivity on a validated test PIMS-TS †† ††

*In the absence of COVID-19

** Excluding superficial thrombophlebitis (including line-associated)

*** G3 or above

**** This will be used as a marker of skewed Th2 responses and will be routinely monitored in participants attending the COVID-19 Pathway and follow-up visits. Only G2 and above.

Brighton collaboration guideline definitions for myocarditis and pericarditis as outlined in appendix D

† In particular, any occurrence of suspected vaccine associated enhanced disease (VAED) as defined by most recent Brighton Collaboration Case Definition (21)

†† In the absence of an infective aetiology (including COVID-19)

†† † As defined in Hy's Law

†† †† The Royal College of Paediatrics and Child Health has outlined a case definition for PIMS-TS.

1. A child presenting with persistent fever, inflammation (neutrophilia, elevated CRP and lymphopaenia) and evidence of single or multi-organ dysfunction (shock, cardiac, respiratory, renal, gastrointestinal or neurological disorder) with additional features (Table 8) This may include children fulfilling full or partial criteria for Kawasaki disease.
2. Exclusion of any other microbial cause, including bacterial sepsis, staphylococcal or streptococcal shock syndromes, infections associated with myocarditis such as enterovirus (waiting for results of these investigations should not delay seeking expert advice).
3. SARS-CoV-2 PCR testing may be positive or negative

AESIs should be collected and recorded in the AE reporting form in REDCap throughout the duration of this study. These should also be reported as SAEs if they fulfil the definition criteria for SAEs. All AESIs not already reported as SAEs should be included in the reports to the DSMB.

11.9.1. Disease enhancement following vaccination

Severe COVID-19 disease will be defined as hospitalisation, with further grading of severity according to the WHO ordinal scale (June 2020).(22) Cases of COVID-19 disease will be examined for the possibility of vaccine associated enhanced disease (VAED). This will be evaluated on the basis of the most recent recommendations of the Brighton Collaboration. Detailed clinical parameters will be collected from medical records and aligned with agreed definitions, as they emerge. Investigations will be defined by the laboratory analysis plan.

11.10. SUSAR reporting

All SUSARs will be reported by the sponsor delegate to the relevant Competent Authority and to the REC and other parties as applicable. For fatal and life-threatening SUSARS, this will be done no later than 7 calendar days after the Sponsor or delegate is first aware of the reaction. Any additional relevant information will be reported within 8 calendar days of the initial report. All other SUSARs will be reported within 15 calendar days.

Principal Investigators will be informed of all SUSARs for the relevant IMP for all studies with the same Sponsor, whether or not the event occurred in the current trial.

11.11. Development Safety Update Reports

A Development Safety Update Report (DSUR) will be prepared annually, reporting on this study, 'Comparing COVID-19 Vaccine Schedule Combinations (Com-COV) (Ethics Ref: 21/SC/0022, IRAS Project ID: 291055) and Comparing COVID-19 Vaccine Schedule Combinations -stage 2 (Com-COV2)(Ethics Ref: 21/SC/0119, IRAS Project ID 297443), within 60 days of the anniversary of the MHRA approval for the 'first' COMCOV study. The DSUR will be submitted by the CI to the Competent Authority, Ethics Committee, HRA (where required), Host NHS Trust and Sponsor.

11.12. Interim reviews

The safety profile will be assessed on an on-going basis by the Investigators. The CI and relevant Investigators (as per the trial delegation log) will also review safety issues and SAEs as they arise.

The DSMB will evaluate safety data as required. The DSMB may also be consulted should safety concerns arise at any point.

11.13. Safety holding rules

There will be no formal pausing rules given the extensive safety database for all vaccines used in this study.

The study can be paused upon advice of the DSMB, Chief Investigator, Study Sponsor, regulatory authority, Ethical Committee(s), for any single event or combination of multiple events which, in their professional opinion, jeopardise the safety of the participants or the reliability of the data.

11.14. Contraception and pregnancy

11.14.1. Contraception

Post-menarchal female participants are required to use an effective form of contraception from enrolment continuously until three months after boost immunisation.

Acceptable forms of contraception for volunteers of female sex include:

- Established use of oral, injected or implanted hormonal methods of contraception
- Placement of an intrauterine device (IUD) or intrauterine system (IUS)
- Total hysterectomy
- Bilateral Tubal Occlusion
- Barrier methods of contraception (condom or occlusive cap with spermicide)
- Male sterilisation, if the vasectomised partner is the sole partner for the subject
- True abstinence, when this is in line with the preferred and usual lifestyle of the subject (Periodic abstinence and withdrawal are not acceptable methods of contraception)

11.14.2. Pregnancy

Should a participant become pregnant during the trial, no further study IMP will be administered. They will be followed up for clinical safety assessment with their ongoing consent and in addition will be followed until pregnancy outcome is determined. We would not routinely perform venepuncture in a pregnant participant unless there is clinical need.

12. STATISTICS

12.1. Sample size

The total sample size of this trial will be 360 with 90 participants in each of the study arms. Since the primary endpoint analysis will be descriptive and therefore no formal sample size calculation is carried out to determine the sample size of the trial. The number has been therefore chosen based on practical constraints. The precisions on estimating the systematic reactions based on 90 per arm are:

True event rate	Precision (normal approximation)	95% exact binomial CI
5%	±4.5%	1.5%-11.7%
10%	±6.2%	4.7%-18.1%

25%	±8.9%	16.5%-35.2%
50%	±10.3%	39.3%-60.7%

For secondary endpoints of immunogenicity, the minimum geometric mean ratios that can be detected with 90% power at two-sided 0.05 based on 90 per arm are:

Standard deviation on log10 scale	Minimum GMR to detect
0.4	1.57 (or 0.64)
0.5	1.75 (or 0.57)
0.6	1.96 (or 0.51)

12.2. Description of statistical methods

A fully detailed statistical analysis plan will be developed and signed by the chief investigator prior to any data analysis being conducted. In brief, the analysis will incorporate the following:

Safety and reactogenicity

All SAEs will be presented for each group using descriptive analyses. Counts and percentages of each local and systemic solicited adverse reaction from diary cards, and all unsolicited AEs, and AEs of special interest will be presented for each group. Comparisons between the arms will be made using Fisher's exact test.

Immunogenicity

Highly skewed ELISA data will be log-transformed prior to analysis. The geometric mean concentration (GMC) and associated 95% confidence interval (CI) will be summarised by computing the anti-log of the mean of the log-transformed data.

The GMRs and CIs post boost will be calculated for each of the heterologous arms when comparing to the homologous arm.

Spike-specific T cell responses (ELISPOT) will be presented as geometric means and confidence intervals, or medians and interquartile ranges if non-normally distributed after log-transformation.

Comparisons of continuous immunogenicity data between different arms or at different time points will be made using a t-test if normal distribution can be rendered after log-transformation. Otherwise, Mann Whitney U test will be used.

12.3. Missing data

Any missing data will be dealt with, if needed, using methods appropriate to the conjectured missing mechanism and level of missing.

12.4. Interim analyses

The analysis on the primary endpoint of reactogenicity, i.e. solicited AEs will be carried out once the 7-day e-diary data become available. The analysis on the secondary endpoint of immunogenicity will be carried out when the D70 immunogenicity data, i.e. 14 days post boost, become available. The interim analyses will provide evidence for policy making as to the recommendation on the second dose of COVID-19 vaccine among adolescents. The interim analyses will be carried out once the data is cleaned and the Study Analysis Plan is signed off. There will be no stopping rule for this interim analysis and the analysis will not affect the continuation of the trial.

After further discussion with Joint Committee on Vaccination and Immunisation (JCVI), the study team decided to carry out another interim analysis in December 2021. In this analysis, we will review the sero-status pre-boost as well as all the available immunogenicity data. These data will be shared with JCVI only to support decision making on boost dose among people aged 12-17. This interim analysis will be descriptive and will not affect the continuation of the trial.

13. DATA MANAGEMENT

The Chief Investigator will be responsible for all data that accrues from the study.

13.1. Access to Data & Data Protection

Direct access will be granted to authorised representatives from the Sponsor, host institution and the regulatory authorities to permit trial-related monitoring, audits and inspections. The study protocol, documentation, data and all other information generated will be held in strict confidence. No information concerning the study or the data will be released to any unauthorised third party, without prior written approval of the sponsor.

13.2. Data Recording

All study data including participant diary will be recorded directly into an EDC system (REDCap) or onto a paper source document for later entry into EDC if direct entry is not available. This includes safety, laboratory and outcome data. Any additional information that needs recording, but is not relevant for the eCRF (e.g signed consent forms) will be recorded on separate paper source documents. All documents will be stored safely and securely in confidential conditions. The EDC online data is stored on University of Oxford servers.

All participant reported adverse event data (both solicited & unsolicited) will be entered onto electronic diary cards (e-diaries) for a maximum of 28 days following administration of the IMP. The e-diary provides a full audit trail of edits and will be reviewed at time-points as indicated in the schedule of events. Any adverse event continuing beyond the period of the diary will be copied into the eCRF as required for safety review.

The participants will be identified by a unique trial specific number and code in any database. The name and any other identifying detail will NOT be included in any trial data electronic file, with the exception of the electronic diaries, for which consent will be obtained to store the participant email address for quality control purposes. Only site research staff and sponsor data managers have access to view the email address.

The EDC system (CRF data) uses a relational database (MySQL/ PostgreSQL) via a secure web interface with data checks applied during data entry to ensure data quality. The database includes a complete suite of features which are compliant with GCP, EU and UK regulations and Sponsor security policies, including a full audit trail, user-based privileges, and integration with the institutional LDAP server. The MySQL and PostgreSQL database and the webserver will both be housed on secure servers maintained by the University of Oxford IT personnel. The servers are in a physically secure location in Europe. Backups will be stored in accordance with the IT department schedule of daily, weekly, and monthly retained for one month, three months, and six months, respectively. The IT servers provide a stable, secure, well-maintained, and high-capacity data storage environment. REDCap is a widely-used, powerful, reliable, well-supported system. Access to the study's database will be restricted to members of the study team by username and password.

13.3. Record keeping

The Investigators will maintain appropriate medical and research records for this trial, in compliance with GCP and regulatory and institutional requirements for the protection of

confidentiality of volunteers. The Chief Investigator, co-Investigators and clinical research nurses will have access to records. The Investigators will permit authorised representatives of the Sponsor(s) and Host institution, as well as ethical and regulatory agencies to examine (and when required by applicable law, to copy) clinical records for the purposes of quality assurance reviews, audits and evaluation of the study safety and progress.

Following completion of the study, identifiable information such as contact details will be stored for a minimum of 5 years and until the youngest participant turns 21 years old. This includes storage of consent forms. Storage of these data will be reviewed every 5 years and files will be confidentially destroyed if storage is no longer required. Considerations at the time of this review will include the value of retaining this information for participant safety (e.g. to inform participants of unexpected safety signals emerging from post-licensing surveillance), as a resource for the participants (e.g. if they wish to check which vaccines they have received in the study) and any regulatory requirements. De-identified research data maybe be stored indefinitely. If volunteers consent to be contacted for future research, a record of this consent will be recorded, retained and stored securely and separately from the research data. If volunteers consent to have their samples stored and used in future research, information about their consent form will be retained and stored securely as per Biobanking procedures and SOP.

13.4. Source data and Case Report Forms (CRFs)

All protocol-required information will be collected in CRFs designed by the Investigator. All source documents will be filed in the participant file. Source documents are original documents, data, and records from which the participant CRF data are obtained. For this study, these will include, but are not limited to, volunteer consent form, blood results, GP response letters, laboratory records, diaries, medical records and correspondence. In the majority of cases, CRF entries will be considered source data as the CRF is the site of the original recording (i.e. there is no other written or electronic record of data). In this study this will include, but is not limited to medical history, medication records, vital signs, physical examination records, urine assessments, adverse event data and details of vaccinations. All source data and participant files will be stored securely.

To prevent withdrawal of a participant due to relocation, if there is a nearby participating site and with the consent of the participant, copies of relevant participant research records (such as ICF, paper source documents) will be transferred to the local site using secure email addresses

such as nhs.net or by password protected sheets. The electronic research data stored on REDCap will also be transferred to the new site. The original records will be retained by the recruiting site.

13.5. Data Quality

Data collection tools will undergo appropriate validation to ensure that data are collected accurately and completely. Datasets provided for analysis will be subject to quality control processes to ensure analysed data is a true reflection of the source data.

Trial data will be managed in compliance with local data management SOPs. If additional, study specific processes are required, an approved Data Management Plan will be implemented

14. Quality Assurance Procedures

14.1. Risk Assessment

The trial will be conducted in accordance with the current approved protocol, GCP, relevant regulations and standard operating procedures. A risk assessment and monitoring plan will be prepared before the study opens and will be reviewed as necessary over the course of the trial to reflect significant changes to the protocol or outcomes of monitoring activities.

14.2. Monitoring

Monitoring will be performed according to Good Clinical Practice (GCP) by an external monitor. Following written SOPs, the monitors will verify that the clinical trial is conducted and data are generated, documented and reported in compliance with the protocol, GCP and the applicable regulatory requirements. The investigator sites will provide direct access to all trial related source data/documents and reports for the purpose of monitoring and auditing by the Sponsor or the Host institution and inspection by local and regulatory authorities

14.3. Trial committees

14.3.1. Trial Steering Committee

A Trial Steering Committee will be formed to oversee the study, and advise the Study Management Committee on key issues of study conduct, including, but not limited to, study pauses due to safety concerns on the advice of the DSMB.

14.3.2. Safety Monitoring Committee

A Data Safety Monitoring Board (DSMB) will be convened. The DSMB will evaluate frequency of events, safety and efficacy data as specified in the DSMB charter. The DSMB will make

recommendations concerning the conduct, continuation or modification of the study for safety reasons to the Trial Steering Committee.

The DSMB will review SAEs or AESIs deemed possibly, probably or definitively related to study interventions. The DSMB will be notified within 24 hours of the Investigators' being aware of their occurrence. The DSMB can recommend placing the study on hold if deemed necessary following a study intervention-related SAE.

14.3.3. Study Management Committee

Consists of the site Investigators and the Laboratory lead for Public Health England.

15. PROTOCOL DEVIATIONS

A trial related deviation is a departure from the ethically approved trial protocol or other trial document or process (e.g. consent process or IMP administration) or from Good Clinical Practice (GCP) or any applicable regulatory requirements. Deviations from the protocol will be documented in a protocol deviation form according to SOP OVC027 and filed in the trial master file.

These will be managed as per SOP OVC027.

16. SERIOUS BREACHES

The Medicines for Human Use (Clinical Trials) Regulations contain a requirement for the notification of "serious breaches" to the MHRA within 7 days of the Sponsor becoming aware of the breach.

A serious breach is defined as "A breach of GCP or the trial protocol which is likely to affect to a significant degree –

- (a) the safety or physical or mental integrity of the subjects of the trial; or
- (b) the scientific value of the trial".

In the event that a serious breach is suspected the Sponsor must be contacted within 1 working day. In collaboration with the CI the serious breach will be reviewed by the Sponsor and, if appropriate, the Sponsor will report it to the REC committee, Regulatory authority and the relevant NHS host organisation within seven calendar days.

17. ETHICAL AND REGULATORY CONSIDERATIONS

17.1. Declaration of Helsinki

The Investigator will ensure that this trial is conducted in accordance with the principles of the Declaration of Helsinki.

17.2. Guidelines for Good Clinical Practice

The Investigator will ensure that this trial is conducted in accordance with relevant regulations and with Good Clinical Practice.

17.3. Approvals

Following Sponsor approval, the protocol, informed consent form, participant information sheet and any proposed advertising material will be submitted to an appropriate Research Ethics Committee (REC), HRA (where required), regulatory authorities (MHRA in the UK), and host institution(s) for written approval. No amendments to this protocol will be made without consultation with, and agreement of, the Sponsor.

The Investigator is responsible for ensuring that changes to an approved trial, during the period for which regulatory and ethical committee(s) approval has already been given, are not initiated without regulatory and ethical committee(s)' review and approval except to eliminate apparent immediate hazards to the subject (i.e. as an Urgent Safety Measure).

17.4. Other Ethical Considerations

First degree family members of the study team are not eligible for inclusion in the trial given the potential for them to be unblinded to the vaccine the participant has received, and for this to influence completion of reactogenicity diaries.

17.5. Reporting

The CI shall submit once a year throughout the clinical trial, or on request, an Annual Progress Report to the REC, HRA (where required), host organisation, funder (where required) and Sponsor. In addition, an End of Trial notification and final report will be submitted to the MHRA, the REC, host organisation and Sponsor.

17.6. Transparency in Research

Prior to the recruitment of the first participant, the trial will have been registered on a publicly accessible database. Results will be uploaded to the European Clinical Trial (EudraCT) Database within 12 months of the end of trial declaration by the CI or their delegate. Where the trial has

been registered on multiple public platforms, the trial information will be kept up to date during the trial, and the CI or their delegate will upload results to all those public registries within 12 months of the end of the trial declaration.

17.7. Participant confidentiality

The study will comply with the UK General Data Protection Regulation (GDPR) and Data Protection Act 2018, which require data to be de-identified as soon as it is practical to do so. The processing of personal data of participants will be minimised by making use of a unique participant study number only on all study documents and any electronic database(s), with the exception of informed consent forms, participant ID log, electronic diaries and any other documents required for participant management. All documents will be stored securely and only accessible by study staff and authorised personnel. The study staff will safeguard the privacy of participants' personal data. A separate confidential file containing identifiable information will be stored in a secured location in accordance with the current data protection legislation.

17.8. Expenses and Benefits

Participants will be offered £10 reimbursement per visit towards travel and expenses. This may be in the form of gift cards/vouchers. Reimbursement may not be given at each visit (for example, we may give a £20 voucher at every other visit). The exact arrangements for reimbursement may vary between sites.

18. FINANCE AND INSURANCE

18.1. Funding

The study is funded by the UK Government through the National Institute for Health Research (NIHR).

18.2. Insurance

The University has a specialist insurance policy in place which would operate in the event of any participant suffering harm as a result of their involvement in the research (Newline Underwriting Management Ltd, at Lloyd's of London). NHS indemnity operates in respect of the clinical treatment that is provided and for the conduct of the research at NHS sites.

18.3. Contractual arrangements

Appropriate contractual arrangements will be put in place with all third parties.

19. PUBLICATION POLICY

The Investigators will be involved in reviewing drafts of the manuscripts, abstracts, press releases and any other publications arising from the study. Data from the study may also be used as part of a thesis for a PhD or MD.

20. DEVELOPMENT OF A NEW PRODUCT / PROCESS OR THE GENERATION OF INTELLECTUAL PROPERTY

Ownership of IP generated by employees of the University vests in the University. The University will ensure appropriate arrangements are in place as regards any new IP arising from the trial.

21. ARCHIVING

Study data may be stored electronically on a secure server, and paper notes will be kept in a key-locked filing cabinet at the site. All essential documents will be retained for a minimum of 9 years (until the youngest participant turns 21) after the study has finished with 5 yearly reviews. The need to store study data for longer in relation to licensing of the vaccine will be subject to ongoing review. For effective vaccines that may be licensed, we may store research data securely at the site at least 15 years after the end of the study, subject to adjustments in clinical trials regulations. De-identified research data may be stored indefinitely, but with 5 yearly review.

General archiving procedures will be conducted in compliance to SOP OVC020 Archiving.

22. REFERENCES:

1. Public Health England. COVID-19 vaccine surveillance report published [Internet]. 2021 [cited 2021 Aug 17]. Available from: <https://www.gov.uk/government/news/covid-19-vaccine-surveillance-report-published>
2. Public Health England. JCVI issues updated advice on COVID-19 vaccination of young people aged 16 to 17 [Internet]. 2021 [cited 2021 Aug 17]. Available from: <https://phe-newsroom.prgloo.com/news/jcvi-issues-updated-advice-on-covid-19-vaccination-of-young-people-aged-16-to-17>
3. A Dionne, F Spirotto, S Chamberlain, AL Baker, AJ Powell, A Prakash et al. Association of Myocarditis With BNT162b2 Messenger RNA COVID-19 Vaccine in a Case Series of Children. JAMA Cardiol [Internet]. 2021;(10 August). Available from: <https://jamanetwork.com/journals/jamacardiology/fullarticle/2783052>

4. N Zhu, D Zhang, W Wang, X Li, B Yang, J Song et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N Engl J Med* [Internet]. 2020;382:727–33. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa2001017>
5. R Lu, X Zhao, J Li, P Niu, B Yang, H Wu et al. Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* [Internet]. 2020;395(10224):565–74. Available from: [https://www.thelancet.com/article/S0140-6736\(20\)30251-8/fulltext](https://www.thelancet.com/article/S0140-6736(20)30251-8/fulltext)
6. WHO. WHO Coronavirus (COVID-19) Dashboard [Internet]. Available from: <https://covid19.who.int/>
7. Li F. Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol* [Internet]. 2016;3:237–61. Available from: <https://www.annualreviews.org/doi/abs/10.1146/annurev-virology-110615-042301>
8. P Zhou, XL Yang, XG Wang et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* [Internet]. 2020;579:270–3. Available from: <https://www.nature.com/articles/s41586-020-2012-7>
9. Department of Health and Social Care. Coronavirus (COVID-19): adult social care guidance Information for the adult social care sector on responding to the coronavirus pandemic. [Internet]. 2020. Available from: <https://www.gov.uk/government/collections/coronavirus-covid-19-social-care-guidance>
10. The Recovery Collaborative Group. Dexamethasone in Hospitalized Patients with Covid-19. *N Engl J Med* [Internet]. 2021;384:693–704. Available from: <https://www.nejm.org/doi/full/10.1056/NEJMoa2021436>
11. JH Beigel, KM Tomashek, L. Dodd, AK. Mehta, BS. Zingman, AC. Kalil et al. Remdesivir for the Treatment of Covid-19 — Final Report. *N Engl J Med* [Internet]. 2020;383:1813–26. Available from: <https://www.nejm.org/doi/full/10.1056/nejmoa2007764>
12. Centers for Disease Control and Prevention. COVID data tracker [Internet]. 2020 [cited 2021 Aug 9]. Available from: <https://covid.cdc.gov/covid-data-tracker/#vaccination-demographics-trends>
13. Centers for Disease Control and Prevention. Selected adverse events recorded after COVID-19 vaccination [Internet]. 2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/safety/adverse-events.html>
14. Pfizer Inc. Pfizer and BioNTech Announce Positive Topline Results From Pivotal Trial of COVID-19 Vaccine In Children 5 to 11 Years [Internet]. 2021 [cited 2021 Oct 18]. Available from: <https://www.pfizer.com/news/press-release/press-release-detail/pfizer-and-biontech-announce-positive-topline-results>
15. PT Heath, EP Galiza, DN Baxter, N Boffito, D Browne, F Burns et al. Safety and Efficacy of NVX-CoV2373 Covid-19 Vaccine. *NEJM* [Internet]. 2021;(June 30). Available from: <https://www.nejm.org/doi/full/10.1056/NEJMoa2107659>
16. Department of Health and Social Care. JCVI statement, September 2021: COVID-19 vaccination of children aged 12 to 15 years [Internet]. 2021 [cited 2021 Sep 3]. Available from: <https://www.gov.uk/government/publications/jcvi-statement-september-2021-covid-19-vaccination-of-children-aged-12-to-15-years>
17. J Flood, J Shingleton, E Bennet, B Walker, Z Amin-Choudhury, G Oligbu et al. Paediatric multisystem inflammatory syndrome temporally associated with SARS-CoV-2 (PIMS-TS): Prospective, national surveillance, United Kingdom and Ireland, 2020. *Lancet* [Internet]. 2021;3. Available from: [https://www.thelancet.com/journals/lanepi/article/PIIS2666-7762\(21\)00052-1/fulltext](https://www.thelancet.com/journals/lanepi/article/PIIS2666-7762(21)00052-1/fulltext)
18. Brighton Collaboration. Myocarditis/Pericarditis Case Definition [Internet]. 2021 [cited 2021 Aug 17]. Available from: <https://brightoncollaboration.us/myocarditis-case-definition-update/>
19. TT Shimabukuro, M Cole, J Su. Reports of Anaphylaxis After Receipt of mRNA COVID-19 Vaccines in the US—December 14, 2020-January 18, 2021. *JAMA Insights* [Internet].

- 2021;325(11):1101–2. Available from:
<https://jamanetwork.com/journals/jama/fullarticle/2776557>
20. Public Health England. Vaccine safety and adverse events following immunisation: the green book, chapter 8. In: The Green Book [Internet]. 2013. Available from:
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/147868/Green-Book-Chapter-8-v4_0.pdf
 21. FM Munoz, JP Cramer, CL Dekker, MZ Dudley, BS Graham, M Gurwith et al. Vaccine-associated enhanced disease: Case definition and guidelines for data collection, analysis, and presentation of immunization safety data. Vaccine [Internet]. 2021;39(22):3053–66. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7901381/>
 22. WHO working group on the clinical characterisation and management of COVID-19 infection. A minimal common outcome measure set for COVID-19 clinical research. Lancet Infect Dis [Internet]. 2020;20(8):e192–7. Available from:
[https://www.thelancet.com/journals/laninf/article/PIIS1473-3099\(20\)30483-7/fulltext](https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(20)30483-7/fulltext)

23. APPENDIX A: SCHEDULE OF PROCEDURES

Study timeline	Screening	D0	D56	D70	D84	D182	D364
Study window		Within 120 days of screening	Day 56–70 post prime vaccine at D0 or Day 56 + post prime vaccination in community	Day 14-17 Post D56	Day 28-42 post D56	Day 170-194 Post prime vaccine	Day 224-379 post prime vaccine
Pre - screening online consent	X*						
Informed consent	X	X					
Medical history	X	X					
Interim medical history		X	X	X	X	X	X
Physical examination (as required)	(X)	(X)	(X)	(X)	(X)	(X)	(X)
Urine test (Pregnancy) (if required)		(X)**	X				
COVID-19 vaccination		(X)**	X				
COVID-19		(X)**	X	X	X	X	X

immunogenicity bloods							
Troponin and NT- proBNP blood test		(X)**	X	X			
Saliva and nasal fluid test (if participant consents)		(X)**	X	X			
Diary review			X	X	X		
SAE/AESI check			X	X	X	X	X

* Pre - screening online consent. Restricted to taking a limited medical history online, contacting the GP for further medical record if required and telephone screening visit(s)

**Procedures performed only if prime dose administered in the study

24. APPENDIX B: AMENDMENT HISTORY

Amendment No.	Protocol Version No.	Date issued	Author(s) of changes	Details of Changes made
MHRA change request during MHRA review	1.1	27 Aug 2021	Hannah Robinson	Section 8.3 and 11.14 updated to specify that only post-menarchal females are required to use contraception during the specified period
1	2.0	10 Sep 2021	Philip de Whalley	<ul style="list-style-type: none"> Sections 2, 3, 5.1, 5.2, 7.3, 10.1, 10.3 and 11.8 updated to change one arm of study from ½ dose Novavax to ½ dose Moderna Section 0 updated to include additional DSMB member Section 9.6 updated to include additional diary information to be collected Sections 3, 6 and 25 updated to include D56 anti-nucleocapsid IgG testing for participants receiving first dose of COVID-19 vaccine in the community Section 22 updated. Reference 15 added Section 17.8 updated to clarify that reimbursement is towards travel and expenses Section 3 updated to remove sentence about even distribution across age range Section 23 updated to change D84 study window

2	2.1	20 Sep 2021	Iason Vichos	<ul style="list-style-type: none"> Section 0 Key Trial Contacts: addition of Dr Philip Connor in the Trial Management Group as the Principal Investigator at Public Health Wales
3	2.2	14 October 2021	Sharon Tonner	<ul style="list-style-type: none"> Section 0 Key Trial Contacts: addition of Dr David Turner in the Trial Management Group as the Principal Investigator at Cripps Health Centre
4	3.0	18 Oct 2021	Eimear Kelly, Philip de Whalley, Xinxue Liu	<ul style="list-style-type: none"> IMP dosing for BNT162b2 booster arm 2 changed from 15 µg (0.15ml) to 10 µg (0.10ml) Addition of interim analysis to be reported to JCVI in December 2021

25. APPENDIX C: BLOOD SAMPLING

Study timeline	D0	D56	D70	D84	D182	D364
COVID-19 vaccination	X	X				
Secondary endpoints	Anti-spike IgG* Anti-nucleocapsid IgG* ELIspot* Troponin & NT-proBNP*	Anti-spike IgG Anti-nucleocapsid IgG** ELIspot Troponin & NT-proBNP	Anti-spike IgG ELIspot Troponin & NT-proBNP	Anti-spike IgG PBMC for storage	Anti-spike IgG Anti-nucleocapsid IgG ELIspot	Anti-spike IgG Anti-nucleocapsid IgG ELIspot
Total volume blood per visit	16	16	16	16	16	16
Total volume by end of study						96 ml

* Procedures performed only if prime dose administered in the study

** Testing performed only if prime dose administered in the community

26. APPENDIX D: DEFINITION OF MYOCARDITIS

Brighton Collaboration criteria provide a clinical definition of myocarditis.(18) They distinguish definitive, probable and possible cases, as follows:

Definitive case:

Histopathologic examination of myocardial tissue (autopsy or endomyocardial biopsy) showed myocardial inflammation

OR Elevated myocardial biomarkers (troponin T and/or troponin I), **AND**

Abnormal Imaging Study **EITHER**

Abnormal Cardiac Magnetic Resonance Study (at least 1 of the findings below):

1. Oedema on T2 weighted study, typically patchy in nature
2. Late gadolinium enhancement on T1 weighted study with an increased enhancement ratio between myocardial and skeletal muscle typically involving at least one non-ischemic regional distribution with recover (myocyte injury)

OR Abnormal Echocardiogram (at least 1 of the findings below):

1. New focal or diffuse left or right ventricular function abnormalities (e.g. decreased ejection fraction)
2. Segmental wall motion abnormalities
3. Global systolic or diastolic function depression/abnormality
4. Ventricular dilation
5. Wall thickness change
6. Intracavitary thrombi

Probable case:

Clinical symptoms

EITHER Cardiac symptoms (at least 1 finding below):

- 1.Acute chest pain or pressure
- 2.Palpitations

3. Dyspnea after exercise, at rest, or lying down
4. Diaphoresis
5. Sudden death

OR Non-specific symptoms (at least 2 findings below):

1. Fatigue
2. Abdominal pain
3. Dizziness/syncope
4. Oedema
5. Cough

OR In infants and young children (at least two finding below):

1. Irritability
2. Vomiting
3. Poor feeding
4. Tachypnoea
5. Lethargy

AND Testing supporting diagnosis (Biomarkers, ECHO and ECG)

EITHER Elevated myocardial biomarkers (at least 1 finding below):

1. Troponin I
2. Troponin T
3. CK myocardial band

OR Echocardiogram (ECHO) abnormalities (at least 1 of the findings below):

1. New focal or diffuse left or right ventricular function abnormalities (e.g. decreased ejection fraction)
2. Segmental wall motion abnormalities
3. Global systolic or diastolic function depression/abnormality
4. Ventricular dilation

5. Wall thickness change
6. Intracavity thrombi

OR Electrocardiogram abnormalities that are new and/or normalise on recovery (at least 1 of the findings below):

1. Paroxysmal or sustained atrial or ventricular arrhythmias (premature atrial or ventricular beats, and/or supraventricular or ventricular tachycardia, interventricular conduction delay, abnormal Q waves, low voltages)
2. AV nodal conduction delays or intraventricular conduction defects (atrioventricular block (grade I-III), new bundle branch block)

AND no alternative diagnosis for symptoms.

Possible case:

EITHER Cardiac symptoms (at least 1 finding below):

1. Acute chest pain or pressure
2. Palpitations
3. Dyspnea after exercise, at rest, or lying down
4. Diaphoresis
5. Sudden death

OR Non-specific symptoms (at least 2 findings below):

1. Fatigue
2. Abdominal pain
3. Dizziness/syncope
4. Oedema
5. Cough

OR In infants and young children (at least two finding below):

1. Irritability
2. Vomiting

3. Poor feeding
4. Tachypnoea
5. Lethargy

AND Biomarkers supporting evidence of infection (at least 1 finding below):

1. Elevated CRP
2. Elevated ESR
3. Elevated D-dimer

AND Non-Specific Electrocardiogram (EKG) Abnormalities that are new and/or normalize on recovery (at least 1 finding below):

1. ST segment or T wave abnormalities (elevation or inversion)
2. PACs and VCs

AND No alternative diagnosis for symptoms